arXiv:1301.3625v2 [math.AG] 17 Jan 2013

Real regulator on K_1 of elliptic surfaces

M. Asakura

Contents

1 Introduction **1**

2 Real Regulator map on $H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$ **2**
- 2.1 $H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$ and (in)decomposable parts . . . 3
- 2.2 Beilinson regulator on indecomposable parts . . . 4
- 2.3 $\mathbb{Q}$-structure on determinant of $H^3_{\mathscr{D}}(X/\mathbb{R}, \mathbb{R}(2))$ . . . 6
- 2.4 $e^{false}_{\mathbb{Q}}$ and $e^{false}_{\mathrm{ind},\mathbb{Q}}$ . . . 7

3 Inhomogeneous de Rham cohomology **9**
- 3.1 Notations . . . 9
- 3.2 Inhomogeneous de Rham cohomology . . . 10

4 Explicit computations of regulator on K_1 of elliptic surfaces **13**
- 4.1 1-Extension of MHS's arising from a multiplicative fiber . . . 13
- 4.2 $\mathbb{E}(\overline{U}, D; \mathbb{Z})$ and $\mathbb{E}(\overline{U}, \mathbb{Z})$ . . . 16
- 4.3 Explicit computation of regulator on $H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$ . . . 18

5 Example : $3y^2 + x^3 + (3x + 4t^l)^2$ **18**
- 5.1 Basic data of X . . . 19
- 5.2 Picard-Fuchs operator and V_{dR} . . . 20
- 5.3 Cycles Δ and Γ . . . 20
- 5.4 Another description of $\int_\Delta$ and $\int_\Gamma$ . . . 23

6 Appendix : Gauss-Manin connection and Picard-Fuchs operator **26**
- 6.1 Explicit formulas . . . 26
- 6.2 Proof of Theorem 6.1 . . . 29
- 6.3 Hodge filtration . . . 32

1 Introduction

Let X be a projective nonsingular variety over the complex number field $\mathbb{C}$. Let $H^i_{\mathscr{M}}(X, \mathbb{Z}(j))$ denotes the motivic cohomology group. It is known that $H^i_{\mathscr{M}}(X, \mathbb{Q}(j))$ is isomorphic to

Quillen's K-group $K_{2j-i}(X)^{(j)}$. By the theory of higher Chern classes, we have the regulator map (higher Chern class map)

$$\mathrm{reg} = \mathrm{reg}_{i,j} : H^i_{\mathscr{M}}(X, \mathbb{Z}(j)) \longrightarrow H^i_{\mathscr{D}}(X, \mathbb{Z}(j))$$

to the Deligne-Beilinson cohomology group ([S]). The purpose of this paper is to give a certain method for computations of the regulator map for $(i, j) = (3, 2)$ (namely K_1) and X an elliptic surface.

The cup-product pairing gives rise to a map $\mathbb{C}^\times \otimes \mathrm{Pic}(X) \cong \mathbb{C}^\times \otimes H^2_{\mathscr{M}}(X, \mathbb{Z}(1)) \to H^3_{\mathscr{M}}(X, \mathbb{Z}(2))$. Its image is called the *decomposable* part, and the cokernel is called the *indecomposable* part. The decomposable part does not affect serious difficulty, while the indecomposable part plays the central role in the study of $H^3_{\mathscr{M}}(X, \mathbb{Z}(2))$. According to [GL], we call an element $\xi \in H^3_{\mathscr{M}}(X, \mathbb{Z}(2))$ *regulator indecomposable* if $\mathrm{reg}(\xi)$ does not lie in the image of $\mathbb{C}^\times \otimes \mathrm{NS}(X)$. Obviously regulator indecomposable elements are indecomposable, but the converse is not true in general. Lewis and Gordon constructed regulator indecomposable elements in case X is a product of 'general' elliptic curves ([GL] Theorem 1). Besides there are a lot of other related works, though I don't catch up all of them. On the other hand, in case that X is defined over a number field, the question is more difficult, and as far as I know there are only a few of such examples ([R] §12 etc.).

The real regulator map $\mathrm{reg}_{3,2}$ can be written in terms of harmonic $(1, 1)$-forms. Then one of the technical difficulties appears from the fact that it is not easy to describe harmonic forms explicitly. The key idea in this paper is to use inhomogeneous de Rham cohomology (§3). This enables us to replace harmonic $(1, 1)$-forms with certain "algebraic" differential forms, so that we are free from the difficult harmonic analysis.

This paper is organized as follows. §2 is a quick review of $H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$ and Beilinson regulator. In §3, we prove some elementary results on inhomogeneous de Rham cohomology of elliptic surface. In §4, we give a method of computations of real regulator on K_1 of elliptic surfaces. In §5 we give an example. In particular we construct regulator indecomposable elements for an elliptic surface defined over $\mathbb{Q}$ (Cor.5.2). §6 is an appendix providing proofs of some basic facts on Gauss-Manin connection and Picard-Fuchs operator.

Acknowledgement. A rough idea was inspired during my visit to the University of Alberta in September 2012, especially when I discussed the paper [CDKL] with Professor James Lewis. I would like to express special thanks to him. I'd also like to thank the university members for their hospitality.

2 Real Regulator map on $H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$

For a regular scheme X, let $Z_i(X) = Z^{\dim X - i}(X)$ be the free abelian group of irreducible subvarieties of Krull dimension i. For an integral scheme X, we denote by η_X the field of rational functions on X. For schemes X and T over a base scheme S, we set $X(T) = \mathrm{Mor}_S(T, X)$. and say $x \in X(T)$ a T-valued point of X. If $T = \mathrm{Spec} R$, then we also write $X(R) = X(\mathrm{Spec} R)$.

2.1 $H^3_{\mathscr{M}}(X,\mathbb{Q}(2))$ and (in)decomposable parts

Let X be a smooth variety over a field K. Let $D \subset X$ be an irreducible divisor, and $\widetilde{D} \to D$ the normalization. Let $j : \widetilde{D} \to D \hookrightarrow X$ be the composition. Then we define $\mathrm{Div}_D(f) := j_*\mathrm{Div}_{\widetilde{D}}(f) \in Z^2(X)$ the push-forward of the Weil divisor on $\widetilde{D}$ by j. Let

$$\partial_1 : \bigoplus_{\mathrm{codim} D=1} \eta_D^\times \longrightarrow Z^2(X), \quad [f,D] \longmapsto \mathrm{Div}_D(f)$$

be a homomorphism where we write

$$[f,D] := (\cdots,1,f,1,\cdots) \in \bigoplus_{\mathrm{codim} D=1} \eta_D^\times, \quad (f \text{ is placed in the } D\text{-component}).$$

Let

$$\partial_2 : K_2^M(\eta_X) \longrightarrow \bigoplus_{\mathrm{codim} D=1} \eta_D^\times$$

$$\partial_2\{f,g\} = \sum_{\mathrm{codim} D=1} \left[(-1)^{\mathrm{ord}_D(f)\mathrm{ord}_D(g)} \frac{f^{\mathrm{ord}_D(g)}}{g^{\mathrm{ord}_D(f)}}|_D, D\right]$$

be the tame symbol. Then it is well-known that there is the canonical isomorphism

$$H^3_{\mathscr{M}}(X,\mathbb{Q}(2)) \cong \left(\frac{\mathrm{Ker}(\bigoplus \eta_D^\times \xrightarrow{\partial_1} Z^2(X))}{\mathrm{Im}(K_2^M(\eta_X) \xrightarrow{\partial_2} \bigoplus \eta_D^\times)}\right) \otimes \mathbb{Q}. \tag{2.1}$$

In this paper we always identify the motivic cohomology group $H^3_{\mathscr{M}}(X,\mathbb{Q}(2))$ with the group in the right hand side of (2.1).

Let L/K be a finite extension. Write $X_L := X \times_K L$. Then there is the obvious map

$$L^\times \otimes Z^1(X_L) \longrightarrow H^3_{\mathscr{M}}(X_L,\mathbb{Q}(2)), \quad \lambda \otimes D \longmapsto [\lambda, D].$$

Let $N_{L/K} : H^3_{\mathscr{M}}(X_L,\mathbb{Q}(2)) \to H^3_{\mathscr{M}}(X,\mathbb{Q}(2))$ be the norm map on motivic cohomology. Then we put

$$H^3_{\mathscr{M}}(X,\mathbb{Q}(2))_{\mathrm{dec}} := \sum_{[L:K]<\infty} N_{L/K}(\mathrm{Im}(L^\times \otimes Z^1(X_L) \to H^3_{\mathscr{M}}(X_L,\mathbb{Q}(2))))$$

and call it the *decomposable part*. We put

$$H^3_{\mathscr{M}}(X,\mathbb{Q}(2))_{\mathrm{ind}} := H^3_{\mathscr{M}}(X,\mathbb{Q}(2))/H^3_{\mathscr{M}}(X,\mathbb{Q}(2))_{\mathrm{dec}}$$

and call it the *indecomposable part*. The indecomposable part plays the central role in the study of $H^3_{\mathscr{M}}(X,\mathbb{Q}(2))$.

2.2 Beilinson regulator on indecomposable parts

Let X be a projective smooth variety over $\mathbb{C}$. We denote by $H^\bullet_B(X,\mathbb{Q}) = H^\bullet_B(X(\mathbb{C}),\mathbb{Q})$ (resp. $H_\bullet(X,\mathbb{Q})$) the Betti cohomology (resp. Betti homology). $H^\bullet_{\mathrm{dR}}$ denotes the de Rham cohomology.

By the theory of universal Chern class, there is the *Beilinson regulator map*

$$\mathrm{reg} = \mathrm{reg}_{\mathbb{Q}} : H^3_{\mathscr{M}}(X,\mathbb{Q}(2)) \longrightarrow H^3_{\mathscr{D}}(X,\mathbb{Q}(2)) \cong \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q}, H^2(X,\mathbb{Q}(2))) \tag{2.2}$$

$$= \frac{H^2_B(X,\mathbb{C})}{F^2 + H^2_B(X,\mathbb{Q}(2))} \tag{2.3}$$

where $H^2(X,\mathbb{Q}(2)) = (H^2_B(X,\mathbb{Q}(2)), F^\bullet H^2_{\mathrm{dR}}(X/\mathbb{C}))$ denotes the Hodge structure (of weight -2) and $\mathrm{Ext}^\bullet_{\mathrm{MHS}}$ denotes the Yoneda extension group of mixed Hodge structures. Put

$$H^2_B(X)_{\mathrm{ind}} := H^2_B(X,\mathbb{Q}(1))/\mathrm{NS}(X)\otimes\mathbb{Q}, \quad H^2_{\mathrm{dR}}(X/\mathbb{C})_{\mathrm{ind}} := H^2_{\mathrm{dR}}(X/\mathbb{C})/\mathrm{NS}(X)\otimes\mathbb{C}.$$

$$H^2(X)_{\mathrm{ind}} := (H^2_B(X)_{\mathrm{ind}}, F^\bullet H^2_{\mathrm{dR}}(X/\mathbb{C})_{\mathrm{ind}}) \tag{2.4}$$

the Hodge structure of weight 0. Then the Beilinson regulator map induces the following commutative diagram

$$\begin{tikzcd}
0 \arrow[d] & 0 \arrow[d] \\
H^3_{\mathscr{M}}(X,\mathbb{Q}(2))_{\mathrm{dec}} \arrow[r] \arrow[d] & \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q}, \mathrm{NS}(X)\otimes\mathbb{Q}(1)) \arrow[d] \\
H^3_{\mathscr{M}}(X,\mathbb{Q}(2)) \arrow[r, "\mathrm{reg}"] \arrow[d] & \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q}, H^2(X,\mathbb{Q}(2))) \arrow[d] \\
H^3_{\mathscr{M}}(X,\mathbb{Q}(2))_{\mathrm{ind}} \arrow[r, "\overline{\mathrm{reg}}"] \arrow[d] & \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q}, H^2(X)_{\mathrm{ind}}\otimes\mathbb{Q}(1)) \arrow[d] \\
0 & 0
\end{tikzcd} \tag{2.5}$$

There is an elementary description of the map $\overline{\mathrm{reg}}$, which we recall here. Let $n = \dim X$. Let $\xi = \sum[f_i, D_i] \in \bigoplus \eta^\times_{D_i}$. Let $j : \widetilde{D}_i \to D_i$ be the normalization. Let $\widetilde{Z}_i \subset \widetilde{D}_i$ be the support of $\mathrm{Div}_{\widetilde{D}_i}(f_i)$. Put

$$\widetilde{Z} := \coprod_i \widetilde{Z}_i \subset \widetilde{D} := \coprod_i \widetilde{D}_i, \quad Z := \bigcup_i j(\widetilde{Z}_i) \subset D := \bigcup_i D_i.$$

Consider a commutative diagram

$$\begin{tikzcd}
 & & H^1(\widetilde{D}-\widetilde{Z},\mathbb{Z}(1)) \arrow[d, "a"] & \\
0 \arrow[r] & H_{2n-3}(\widetilde{D},\mathbb{Z}(2-n)) \arrow[r] \arrow[d, "j_*"] & H_{2n-3}(\widetilde{D},\widetilde{Z};\mathbb{Z}(2-n)) \arrow[r, "\delta_1"] \arrow[d, "j_*"] & H_{2n-4}(\widetilde{Z},\mathbb{Z}(2-n)) \arrow[d, "j_*"] \\
0 \arrow[r] & H_{2n-3}(D,\mathbb{Z}(2-n)) \arrow[r, "b"] & H_{2n-3}(D,Z;\mathbb{Z}(2-n)) \arrow[r, "\delta_2"] & H_{2n-4}(Z,\mathbb{Z}(2-n))
\end{tikzcd}$$

with exact rows. Let

$$\nu := \left(\frac{df_i}{f_i}\right) \in H^1(\widetilde{D}-\widetilde{Z},\mathbb{Z}(1)).$$

Suppose $\partial_1(\xi)=0$. This is equivalent to saying $j_*\delta_1 a(\nu)=0$. Therefore there is a unique element $\nu_\xi \in H_{2n-3}(D,\mathbb{Z}(2-n))$ such that $b(\nu_\xi)=j_*a(\nu)$. Note that ν_ξ belongs to the Hodge (0,0)-part because so does ν. Let us consider the localization exact sequence

$$\cdots \longrightarrow H_{2n-2}(X,D;\mathbb{Q}(2-n)) \longrightarrow H_{2n-3}(D,\mathbb{Q}(2-n)) \overset{\delta}{\longrightarrow} H_{2n-3}(X,\mathbb{Q}(2-n)) \longrightarrow \cdots .$$

Since the weight of $H_{2n-3}(X,\mathbb{Q}(2-n))$ is -1, the Hodge $(0,0)$-part of $H_{2n-3}(D,\mathbb{Q}(2-n))$ is contained in the kernel of δ. Put

$$H(n,X,D) := H_{2n-2}(X,\mathbb{Q}(2-n))/H_{2n-2}(D,\mathbb{Q}(2-n)) \overset{\cong}{\longleftarrow} H^2_B(X,\mathbb{Q}(2))/\langle D_i\rangle$$

where $\langle D_i\rangle$ denotes the subgroup generated by the cycle classes of D_i's and the isomorphism is the Poincare duality. Now an exact sequence

$$\begin{tikzcd}
0 \arrow[r] & H(n,X,D) \arrow[r] & H_{2n-2}(X,D;\mathbb{Q}(2-n)) \arrow[r] & \mathrm{Ker}(\delta) \arrow[r] & 0
\end{tikzcd}$$

yields maps

$$\begin{aligned}
\nu_\xi \in \mathrm{Hom}_{\mathrm{MHS}}(\mathbb{Q},H_{2n-3}(D,\mathbb{Q}(2-n)) &\overset{\subset}{\longrightarrow} \mathrm{Hom}_{\mathrm{MHS}}(\mathbb{Q},\mathrm{Ker}(\delta)) \\
&\longrightarrow \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q},H(n,X,D)) \\
&\overset{\cong}{\longleftarrow} \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q},H^2_B(X,\mathbb{Q}(2))/\langle D_i\rangle) \\
&\longrightarrow \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q},H^2(X)_{\mathrm{ind}}\otimes\mathbb{Q}(1)).
\end{aligned}$$

We denote the composition by ρ. Then the following is well-known to specialists, proven by using the Riemann-Roch theorem without denominators ([G], see also [AS] Thm. 11.2).

Theorem 2.1 $\overline{\mathrm{reg}}(\xi)=\pm\rho(\nu_\xi)$. *In more down-to-earth manner, the regulator map is described in the following way. Letting* $\mathrm{NS}^\perp \subset H^{2n-2}_{\mathrm{dR}}(X/\mathbb{C})$ *be the exact annihilator of* $\mathrm{NS}(X)$*, there is an isomorphism*

$$\mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q},H^2(X)_{\mathrm{ind}}\otimes\mathbb{Q}(1)) \cong \mathrm{Coker}[H_{2n-2}(X,\mathbb{Q}(2-n)) \overset{\Phi}{\to} \mathrm{Hom}(F^{2n-3}\mathrm{NS}^\perp,\mathbb{C})]$$

where

$$\Phi(\Delta) = \left[\omega \longmapsto \int_{\Delta} \omega\right].$$

Let $\Gamma \in H_{2n-2}(X, D; \mathbb{Q}(2-n))$ *be an arbitrary lifting of* γ*. Then we have*

$$\overline{\mathrm{reg}}(\xi) = \left[\omega \longmapsto \int_{\Gamma} \omega\right]$$

under the above isomorphism.

The *real regulator map* is the composition of $\mathrm{reg}_{\mathbb{Q}}$ and the canonical map

$$\mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{Q}, H^2(X, \mathbb{Q}(2))) \longrightarrow \mathrm{Ext}^1_{\mathbb{R}\text{-}\mathrm{MHS}}(\mathbb{R}, H^2(X, \mathbb{R}(2)))$$

to the extension group of real mixed Hodge structures, which we denote by $\mathrm{reg}_{\mathbb{R}}$ or $\overline{\mathrm{reg}}_{\mathbb{R}}$.

$$\mathrm{reg}_{\mathbb{R}} : H^3_{\mathscr{M}}(X, \mathbb{Q}(2)) \longrightarrow \mathrm{Ext}^1_{\mathbb{R}\text{-}\mathrm{MHS}}(\mathbb{R}, H^2(X, \mathbb{R}(2))) \cong H^2_B(X, \mathbb{R}(1)) \cap H^{1,1}, \tag{2.6}$$

$$\overline{\mathrm{reg}}_{\mathbb{R}} : H^3_{\mathscr{M}}(X, \mathbb{Q}(2)) \to \mathrm{Ext}^1_{\mathbb{R}\text{-}\mathrm{MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1)) \cong (H^2_B(X)_{\mathrm{ind}} \otimes \mathbb{R}) \cap H^{1,1}. \tag{2.7}$$

2.3 $\mathbb{Q}$-structure on determinant of $H^3_{\mathscr{D}}(X/\mathbb{R}, \mathbb{R}(2))$

Suppose that X is a projective smooth variety over $\mathbb{Q}$. Write $X_{\mathbb{C}} := X \times_{\mathbb{Q}} \mathbb{C}$. The *infinite Frobenius* map F_∞ is defined to be the anti-holomorphic map on $X(\mathbb{C}) = \mathrm{Mor}_{\mathbb{Q}}(\mathrm{Spec}\mathbb{C}, X)$ induced from the complex conjugation on $\mathrm{Spec}\mathbb{C}$. For a subring $A \subset \mathbb{R}$, the infinite Frobenius map acts on the Deligne-Beilinson complex $A_X(j)_{\mathscr{D}}$ in a canonical way, so that we have the involution on $H^\bullet_{\mathscr{D}}(X_{\mathbb{C}}, A(j))$, which we denote by the same notation F_∞. We define

$$H^\bullet_{\mathscr{D}}(X/\mathbb{R}, A(j)) := H^\bullet_{\mathscr{D}}(X_{\mathbb{C}}, A(j))^{F_\infty = 1}$$

the fixed part by F_∞. We call it the *real Deligne-Beilinson cohomology*. Since the action of F_∞ is compatible via the Beilinson regulator map, we have

$$\begin{aligned} \mathrm{reg}_{\mathbb{R}} : H^3_{\mathscr{M}}(X, \mathbb{Q}(2)) \longrightarrow H_{\mathscr{D}} :=& \mathrm{Ext}^1_{\mathbb{R}\text{-}\mathrm{MHS}}(\mathbb{R}, H^2(X_{\mathbb{C}}, \mathbb{R}(2)))^{F_\infty=1} && (2.8)\\ \cong& \frac{H^2_B(X_{\mathbb{C}}, \mathbb{R}(1))^{F_\infty=1}}{F^2 H^2_{\mathrm{dR}}(X/\mathbb{R})}, && (2.9) \end{aligned}$$

and

$$\begin{aligned} \overline{\mathrm{reg}}_{\mathbb{R}} : H^3_{\mathscr{M}}(X, \mathbb{Q}(2)) \longrightarrow H_{\mathscr{D},\mathrm{ind}} :=& \mathrm{Ext}^1_{\mathbb{R}\text{-}\mathrm{MHS}}(\mathbb{R}, H^2(X_{\mathbb{C}})_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty=1} && (2.10)\\ \cong& \frac{[H^2_B(X_{\mathbb{C}})_{\mathrm{ind}} \otimes \mathbb{R}]^{F_\infty=1}}{F^2 H^2_{\mathrm{dR}}(X/\mathbb{R})}. && (2.11) \end{aligned}$$

There are the canonical $\mathbb{Q}$-structures $e_{\mathbb{Q}}$ and $e_{\mathrm{ind},\mathbb{Q}}$ on the determinant vector spaces $\det H_{\mathscr{D}}$ and $\det H_{\mathscr{D},\mathrm{ind}}$:

$$\mathbb{R} \cdot e_{\mathbb{Q}} = \det H_{\mathscr{D}}, \quad \mathbb{R} \cdot e_{\mathrm{ind},\mathbb{Q}} = \det H_{\mathscr{D},\mathrm{ind}}.$$

Here we recall the definition. The isomorphisms (2.9) and (2.11) induce

$$\det H_{\mathscr{D}} \cong \det[H_B^2(X_{\mathbb{C}}, \mathbb{R}(1))^{F_\infty=1}] \otimes [\det F^2 H^2_{\mathrm{dR}}(X/\mathbb{R})]^{-1}, \tag{2.12}$$

and

$$\det H_{\mathscr{D},\mathrm{ind}} \cong \det[(H_B^2(X_{\mathbb{C}})_{\mathrm{ind}} \otimes \mathbb{R})^{F_\infty=1}] \otimes [\det F^2 H^2_{\mathrm{dR}}(X/\mathbb{R})]^{-1}. \tag{2.13}$$

The right hand sides of (2.12) and (2.13) have the $\mathbb{Q}$-structures induced from the $\mathbb{Q}$-structures

$$H_B^2(X_{\mathbb{C}}, \mathbb{Q}(1))^{F_\infty=1}, \quad H_B^2(X_{\mathbb{C}})_{\mathrm{ind}}^{F_\infty=1}, \quad F^2 H^2_{\mathrm{dR}}(X/\mathbb{Q}).$$

The $\mathbb{Q}$-structures $e_{\mathbb{Q}}$ and $e_{\mathrm{ind},\mathbb{Q}}$ are defined to be the corresponding one:

$$\mathbb{Q} \cdot e_{\mathbb{Q}} \cong \det[H_B^2(X_{\mathbb{C}}, \mathbb{Q}(1))^{F_\infty=1}] \otimes [\det F^2 H^2_{\mathrm{dR}}(X/\mathbb{Q})]^{-1}, \tag{2.14}$$

$$\mathbb{Q} \cdot e_{\mathbb{Q},\mathrm{ind}} \cong \det[H_B^2(X_{\mathbb{C}})_{\mathrm{ind}}^{F_\infty=1}] \otimes [\det F^2 H^2_{\mathrm{dR}}(X/\mathbb{Q})]^{-1}. \tag{2.15}$$

2.4 $e_{\mathbb{Q}}^{\mathit{false}}$ and $e_{\mathrm{ind},\mathbb{Q}}^{\mathit{false}}$

We introduce other $\mathbb{Q}$-structures $e_{\mathbb{Q}}^{\mathit{false}}$ and $e_{\mathrm{ind},\mathbb{Q}}^{\mathit{false}}$ on $\det H_{\mathscr{D}}$ and $\det H_{\mathscr{D},\mathrm{ind}}$. For simplicity, we assume $\dim X = 2$. Put

$$H_2(X_{\mathbb{C}}, \mathbb{Q})_{\mathrm{ind}} := H_2(X_{\mathbb{C}}, \mathbb{Q})/(\mathrm{NS}(X_{\mathbb{C}}) \otimes \mathbb{Q}(1)) \cong H_B^2(X_{\mathbb{C}})_{\mathrm{ind}} \otimes \mathbb{Q}(1),$$

$$H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}} := \mathrm{Coim}(H^2_{\mathrm{dR}}(X/\mathbb{Q}) \longrightarrow H^2_{\mathrm{dR}}(X/\mathbb{C})/(\mathrm{NS}(X_{\mathbb{C}}) \otimes \mathbb{C})).$$

Note that $H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}} \otimes \mathbb{C}$ coincides with "$H^2_{\mathrm{dR}}(X_{\mathbb{C}}/\mathbb{C})_{\mathrm{ind}}$" in (2.4). There are exact sequences

$$0 \longrightarrow H_2(X_{\mathbb{C}}, \mathbb{R})^{F_\infty=1} \longrightarrow \mathrm{Hom}(F^1 H^2_{\mathrm{dR}}(X/\mathbb{Q}), \mathbb{R}) \longrightarrow H_{\mathscr{D}} \longrightarrow 0 \tag{2.16}$$

$$0 \longrightarrow H_2(X_{\mathbb{C}}, \mathbb{R})_{\mathrm{ind}}^{F_\infty=1} \longrightarrow \mathrm{Hom}(F^1 H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}}, \mathbb{R}) \longrightarrow H_{\mathscr{D},\mathrm{ind}} \longrightarrow 0 \tag{2.17}$$

under the canonical isomorphisms

$$H_B^2(X_{\mathbb{C}}, \mathbb{C}) \cong H^2_{\mathrm{dR}}(X/\mathbb{C}), \quad H_B^2(X_{\mathbb{C}}, \mathbb{Q}(2)) \cong H_2(X_{\mathbb{C}}, \mathbb{Q}). \tag{2.18}$$

Then the $\mathbb{Q}$-structures

$$H_2(X_{\mathbb{C}}, \mathbb{Q})^{F_\infty=1}, \quad H_2(X_{\mathbb{C}}, \mathbb{R})_{\mathrm{ind}}^{F_\infty=1}, \quad H^2_{\mathrm{dR}}(X/\mathbb{Q}), \quad H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}},$$

induce $e_{\mathbb{Q}}^{\mathit{false}}$ and $e_{\mathrm{ind},\mathbb{Q}}^{\mathit{false}}$:

$$\mathbb{Q} \cdot e_{\mathbb{Q}}^{\mathit{false}} \cong [\det H_2(X_{\mathbb{C}}, \mathbb{Q})^{F_\infty=1}]^{-1} \otimes [\det F^1 H^2_{\mathrm{dR}}(X/\mathbb{Q})]^{-1}, \tag{2.19}$$

$$\mathbb{Q} \cdot e_{\mathrm{ind},\mathbb{Q}}^{\mathit{false}} \cong [\det H_2(X_{\mathbb{C}}, \mathbb{Q})_{\mathrm{ind}}^{F_\infty=1}]^{-1} \otimes [\det F^1 H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}}]^{-1}. \tag{2.20}$$

Proposition 2.2 *Let us put*

$$r := \dim H_2(X_{\mathbb{C}}, \mathbb{Q})^{F_\infty=1} = \dim H^2_B(X_{\mathbb{C}}, \mathbb{Q}(1))^{F_\infty=-1},$$

$$s := \dim H_2(X_{\mathbb{C}}, \mathbb{Q})^{F_\infty=1}_{\text{ind}} = \dim H^2_B(X_{\mathbb{C}})^{F_\infty=-1}_{\text{ind}} = r - \dim \mathrm{NS}(X_{\mathbb{C}})^{F_\infty=-1}.$$

Then

$$\mathbb{Q} \cdot e^{\textit{false}}_{\mathbb{Q}} = \mathbb{Q} \cdot e_{\mathbb{Q}} \otimes \mathbb{Q}(-r) \otimes \det H^2_{\mathrm{dR}}(X/\mathbb{Q}) \otimes [\det H^2_B(X_{\mathbb{C}}, \mathbb{Q}(1))]^{-1},$$

$$\mathbb{Q} \cdot e^{\textit{false}}_{\text{ind},\mathbb{Q}} = \mathbb{Q} \cdot e_{\text{ind},\mathbb{Q}} \otimes \mathbb{Q}(-s) \otimes \det H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\text{ind}} \otimes [\det H^2_B(X_{\mathbb{C}})_{\text{ind}}]^{-1},$$

where we mean

$$\det H^2_{\mathrm{dR}}(X/\mathbb{Q}) \otimes (\det H^2_B(X_{\mathbb{C}}, \mathbb{Q}(1)))^{-1} \subset \det H^2_{\mathrm{dR}}(X/\mathbb{C}) \otimes [\det H^2_B(X_{\mathbb{C}}, \mathbb{C})]^{-1} \overset{(2.18)}{\cong} \mathbb{C},\ \textit{etc.}$$

Proof. Write $H_B := H^2_B(X_{\mathbb{C}}, \mathbb{Q}(1))$, $H_{B,\text{ind}} := H^2_B(X_{\mathbb{C}})_{\text{ind}}$, $F^\bullet H_{\mathrm{dR}} = F^\bullet H^2_{\mathrm{dR}}(X/\mathbb{Q})$ and $F^\bullet H_{\mathrm{dR,ind}} = F^\bullet H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\text{ind}}$ simply. By the Poincare duality,

$$\det F^1 H_{\mathrm{dR}} = [\det H_{\mathrm{dR}}]^{-1} \otimes \det F^2 H_{\mathrm{dR}}, \quad \det F^1 H_{\mathrm{dR,ind}} = [\det H_{\mathrm{dR,ind}}]^{-1} \otimes \det F^2 H_{\mathrm{dR}}.$$

Moreover

$$\det[H_2(X_{\mathbb{C}}, \mathbb{Q})^{F_\infty=1}] = \det[H^2_B(X_{\mathbb{C}}, \mathbb{Q}(2))^{F_\infty=1}] = \det[H_B^{F_\infty=-1} \otimes \mathbb{Q}(1)] = \mathbb{Q}(r) \otimes \det H_B^{F_\infty=-1}$$

and

$$\det[H_2(X_{\mathbb{C}}, \mathbb{Q})_{\text{ind}}]^{F_\infty=1} = \det[H^{F_\infty=-1}_{B,\text{ind}} \otimes \mathbb{Q}(1)] = \mathbb{Q}(s) \otimes \det H^{F_\infty=-1}_{B,\text{ind}}.$$

Therefore we have

$$\begin{aligned}\mathbb{Q} \cdot e^{\textit{false}}_{\mathbb{Q}} \otimes e^{-1}_{\mathbb{Q}} &= \mathbb{Q}(-r) \otimes [\det H_B^{F_\infty=-1}]^{-1} \otimes [\det H_B^{F_\infty=1}]^{-1} \otimes [\det H_{\mathrm{dR}}] \\ &= \mathbb{Q}(-r) \otimes [\det H_B]^{-1} \otimes [\det H_{\mathrm{dR}}]\end{aligned}$$

by (2.14) and (2.19), and

$$\begin{aligned}\mathbb{Q} \cdot e^{\textit{false}}_{\text{ind},\mathbb{Q}} \otimes e^{-1}_{\text{ind},\mathbb{Q}} &= \mathbb{Q}(-s) \otimes [\det H^{F_\infty=-1}_{B,\text{ind}}]^{-1} \otimes [\det H^{F_\infty=1}_{B,\text{ind}}]^{-1} \otimes [\det H_{\mathrm{dR,ind}}] \\ &= \mathbb{Q}(-s) \otimes [\det H_{B,\text{ind}}]^{-1} \otimes [\det H_{\mathrm{dR,ind}}]\end{aligned}$$

by (2.15) and (2.20). This completes the proof. □

Remark 2.3 *The Poincare duality implies*

$$(\det H_B)^{\otimes 2} \cong H^4_B(X_{\mathbb{C}}, \mathbb{Q}(2))^{\otimes m} \cong H^4_{\mathrm{dR}}(X/\mathbb{Q})^{\otimes m} \cong (\det H_{\mathrm{dR}})^{\otimes 2},$$

and

$$(\det H_{B,\text{ind}})^{\otimes 2} \cong H^4_B(X_{\mathbb{C}}, \mathbb{Q}(2))^{\otimes m'} \cong H^4_{\mathrm{dR}}(X/\mathbb{Q})^{\otimes m'} \cong (\det H_{\mathrm{dR,ind}})^{\otimes 2}.$$

Therefore $(\det H_{\mathrm{dR}} \otimes [\det H_B]^{-1})$ *and* $(\det H_{\mathrm{dR,ind}} \otimes [\det H_{B,\text{ind}}]^{-1})$ *are contained in* $\sqrt{\mathbb{Q}^\times}$ *(possibly rational numbers).*

3 Inhomogeneous de Rham cohomology

3.1 Notations

Let K be a field of characteristic 0. Let $f : X \to C$ be an elliptic surface with a section $e : C \to X$. This means that X (resp. C) is a projective smooth surface (resp. curve) over K, and the generic fiber of f is an elliptic curve. Hereafter we assume that the j-invariant of f is not constant, namely, f is not isotrivial.

Throughout §3 and §4 we use the following notations.

- $D \subset X$ is the sum of the multiplicative fibers. Put $T_1 = f(D) \subset C$.
- $E \subset X$ is the sum of the additive fibers. Put $T_2 = f(E) \subset C$.
- $T = T_1 + T_2$, $S = C - T$ and $\overline{S} = C - T_2$.
- $U = X - (E + D)$ and $\overline{U} = X - E$.
- Let $F \subset S$ be the locus satisfying the following. Around a neighborhood of $s \in S$, f is written by a Weierstrass form $y^2 = 4x^3 - g_2 x - g_3$ $(g_2, g_3 \in \mathscr{O}_{S,s})$. Let $j = 1728 g_2^3/(g_2^3 - 27 g_3^2)$ be the j-invariant. Then $s \in F$ if and only if

  $$\frac{g_2}{g_3}\frac{dj}{j} \in \mathscr{O}_{S,s} \otimes \Omega^1_S \cong \mathscr{O}_{S,s} \quad (\Omega^1_S := \Omega^1_{S/K})$$

  is a $\mathscr{O}_{S,s}$-base (cf. Prop. 6.3).
- $U^o := U - f^{-1}(F)$ and $S^o := S - F$.
- Write $X_{\overline{K}} = X \times_K \overline{K}$. Let $\mathrm{NF}(X_{\overline{K}}) \subset \mathrm{NS}(X_{\overline{K}})$ denotes the subgroup of the Neron-Severi group generated by $e(C)$ and irreducible components of $D_{\overline{K}} + E_{\overline{K}}$.
- $\mathrm{NF}_{\mathrm{dR}}(X) = H^2_{\mathrm{dR}}(X/K) \cap (\mathrm{NF}(X_{\overline{K}}) \otimes_{\mathbb{Z}} \overline{K}) \subset H^2_{\mathrm{dR}}(X_{\overline{K}}/\overline{K})$.
- $V_{\mathrm{dR}} = \mathrm{Im}(\mathit{\Gamma}(\overline{U}, \Omega^2_{\overline{U}}) \to H^2_{\mathrm{dR}}(\overline{U}/K))$.
- $V_{\mathrm{PF}} = \mathrm{Im}(\mathit{\Gamma}(U^o, \Omega^2_{U^o}) \to H^2_{\mathrm{dR}}(U^o/K))$.

$$\begin{tikzcd}
U^o \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & U \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & \overline{U} \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & X \arrow[d, "f"] \\
S^o \arrow[r] & S \arrow[r] & \overline{S} \arrow[r] & C
\end{tikzcd}$$

Remark 3.1 *The intersection pairing* $\mathrm{NF}(X_{\overline{K}}) \otimes \mathrm{NF}(X_{\overline{K}}) \otimes \mathbb{Q} \to \mathbb{Q}$ *is non-degenerate. This is proven on a case-by-case analysis by using the classification of singular fibers (e.g. [Si] IV, Thm.8.2)*

Remark 3.2 $\mathrm{NF}_{\mathrm{dR}}(X) \otimes_K \overline{K} = \mathrm{NF}(X_{\overline{K}}) \otimes_{\mathbb{Z}} \overline{K}$ *in* $H^2_{\mathrm{dR}}(X_{\overline{K}}/\overline{K})$. *This is proven by using [AEC] II Lemma 5.8.1.*

3.2 Inhomogeneous de Rham cohomology

We put $\mathscr{H} := R^1 f_* \Omega^\bullet_{U/S}$ and

$$\mathscr{H}^{1,0} := f_* \Omega^1_{U/S}, \quad \mathscr{H}^{0,1} := \mathscr{H}/\mathscr{H}^{1,0} \cong R^1 f_* \mathscr{O}_U.$$

Let $\nabla : \mathscr{H} \to \mathscr{H} \otimes \Omega^1_S$ be the Gauss-Manin connection. We call

$$H^q_{\mathrm{dR}}(S^o, \mathscr{H}) := H^q_{\mathrm{zar}}(S^o, \mathscr{H} \xrightarrow{\nabla} \mathscr{H} \otimes \Omega^1_S)$$

the *inhomogeneous de Rham cohomology* of X. An exact sequence

$$0 \longrightarrow \Omega^{\bullet-1}_{U^o/S^o} \otimes \Omega^1_{S^o} \longrightarrow \Omega^\bullet_{U^o} \longrightarrow \Omega^\bullet_{U^o/S^o} \longrightarrow 0$$

yields an exact sequence

$$0 \longrightarrow H^1_{\mathrm{dR}}(S^o, \mathscr{H}) \longrightarrow H^2_{\mathrm{dR}}(U^o/K) \longrightarrow H^2_{\mathrm{dR}}(U^o/S^o) \xrightarrow{\mathrm{id} \otimes d} H^2_{\mathrm{dR}}(U^o/S^o) \otimes \Omega^1_{S^o} \tag{3.1}$$

where $d : \mathscr{O}(S^o) \to \Omega^1_{S^o}$ is the differential operator given by $f \mapsto df$ (we use the same notation "$\Omega^1_{S^o}$" for $\Gamma(S^o, \Omega^1_{S^o})$ since it will be clear from the context which is meant). Since the characteristic of K is 0, the kernel of $(\mathrm{id} \otimes d)$ is a one-dimensional K-vector space generated by the cycle class of the section $e(C)$.

Proposition 3.3 *The inhomogeneous de Rham cohomology is described by the Picard-Fuchs operator:*

$$H^1_{\mathrm{dR}}(S^o, \mathscr{H}) \cong \mathrm{Coker}(\Gamma(S^o, \mathscr{H}^{1,0}) \xrightarrow{PF} \Gamma(S^o, \mathscr{H}^{0,1} \otimes \Omega^1_{S^o})) \cong V_{\mathrm{PF}}.$$

Proof. The first isomorphism follows from (6.8) in Appendix. Since $\Gamma(S^o, \mathscr{H}^{0,1} \otimes \Omega^1_{S^o}) = \Gamma(U^o, \Omega^2_{U^o})$, the first isomorphism yields $H^1_{\mathrm{dR}}(S^o, \mathscr{H}) = \mathrm{Im}\Gamma(U^o, \Omega^2_{U^o}) = V_{\mathrm{PF}}$. $\square$

Proposition 3.4 *Assume $E \neq \emptyset$ (hence $\overline{S}$ is affine).*

1. *Let $\mathrm{NF}_{\mathrm{dR}}(X)^\perp$ be the orthogonal complements of $\mathrm{NF}_{\mathrm{dR}}(X)$ in $H^2_{\mathrm{dR}}(X/K)$ with respect to the cup-product pairing. Then $\mathrm{NF}_{\mathrm{dR}}(X)^\perp \overset{\cong}{\to} V_{\mathrm{dR}}$.*
2. $$0 \longrightarrow V_{\mathrm{dR}} \longrightarrow V_{\mathrm{PF}} \longrightarrow H^{\mathrm{dR}}_1(D/K) \oplus H^{\mathrm{dR}}_1(f^{-1}(F)/K) \longrightarrow 0$$
3. *Let $\langle e(C) \rangle_K \subset H^2_{\mathrm{dR}}(X/K)$ denotes the sub K-module generated by the cycle class of $e(C)$. Then $V_{\mathrm{PF}} \oplus \langle e(C) \rangle_K \overset{\cong}{\to} H^2_{\mathrm{dR}}(U^o/K)$.*
4. *Let $\langle e(C), \widetilde{D} \rangle_{\overline{K}} \subset H^2_{\mathrm{dR}}(X_{\overline{K}}/\overline{K})$ denotes the sub $\overline{K}$-module generated by the cycle classes of $e(C)$ and irreducible components of $D_{\overline{K}}$. Put $\langle e(C), \widetilde{D} \rangle_K := \langle e(C), \widetilde{D} \rangle_{\overline{K}} \cap H^2_{\mathrm{dR}}(X/K)$. Then $V_{\mathrm{dR}} \oplus \mathrm{Im}\langle e(C), \widetilde{D} \rangle_K \overset{\cong}{\to} H^2_{\mathrm{dR}}(\overline{U}/K)$ where $\mathrm{Im}\langle e(C), \widetilde{D} \rangle_K$ denotes the image in $H^2_{\mathrm{dR}}(\overline{U}/K)$.*

(5)

$$0 \longrightarrow V_{\mathrm{dR}} \longrightarrow H^2_{\mathrm{dR}}(\overline{U}/K) \longrightarrow H^2_{\mathrm{dR}}(D/K).$$

Proof. Note that $\mathrm{NF}_{\mathrm{dR}}(X)^{\perp}\otimes_K\overline{K} = (\mathrm{NF}(X_{\overline{K}})\otimes_{\mathbb{Z}}\overline{K})^{\perp}$ by Rem. 3.2, and $\langle e(C), D\rangle_K\otimes\overline{K} = \langle e(C), D\rangle_{\overline{K}}$ as well. Therefore we may assume $K = \overline{K}$ throughout the proof.

We first show (3). Let us consider the spectral sequence

$$E_1^{pq} = H^q(U^o, \Omega^p_{U^o}) \Longrightarrow H^{p+q}_{\mathrm{dR}}(U^o/K).$$

Since S^o is affine, $E_1^{pq} = H^q(U^o, \Omega^p_{U^o}) = H^0(S^o, R^qf_*\Omega^p_{U^o})$ and this vanishes unless $p \leq 2$ and $q \leq 1$. Therefore we have

$$E_1^{20} = \mathit{\Gamma}(U^o, \Omega^2_{U^o}) \xrightarrow{\text{surj.}} E_3^{20} = E_\infty^{20}, \quad E_2^{11} = E_\infty^{11}, \quad E_2^{02} = 0$$

and hence an exact sequence

$$0 \longrightarrow V_{\mathrm{PF}} \longrightarrow H^2_{\mathrm{dR}}(U^o/K) \longrightarrow E_2^{11} \longrightarrow 0.$$

Thus it is enough to show $E_2^{11} = \langle e(C)\rangle_K$. The exact sequence

$$0 \longrightarrow f^*\Omega^1_{S^o} \longrightarrow \Omega^1_{U^o} \longrightarrow \Omega^1_{U^o/S^o} \longrightarrow 0$$

yields

$$0 \to \Omega^1_{S^o} \to f_*\Omega^1_{U^o} \to f_*\Omega^1_{U^o/S^o} \xrightarrow{\overline{\nabla}} \Omega^1_{S^o} \otimes R^1f_*\mathscr{O}_{U^o} \to R^1f_*\Omega^1_{U^o} \to R^1f_*\Omega^1_{U^o/S^o} \to 0.$$

By Prop. 6.3 one has

$$\Omega^1_{S^o} \xrightarrow{\cong} f_*\Omega^1_{U^o}, \quad R^1f_*\Omega^1_{U^o} \xrightarrow{\cong} R^1f_*\Omega^1_{U^o/S^o}.$$

Hence

$$E_1^{11} = H^0(S^o, R^1f_*\Omega^1_{U^o}) = H^0(S^o, R^1f_*\Omega^1_{U^o/S^o}) = \mathit{\Gamma}(S^o, \mathscr{O}_{S^o}) \otimes [e(C)],$$

$$E_1^{21} = H^0(S^o, R^1f_*\Omega^2_{U^o}) = H^0(S^o, \Omega^1_{S^o} \otimes R^1f_*\Omega^1_{U^o/S^o}) = \mathit{\Gamma}(S^o, \Omega^1_{S^o}) \otimes [e(C)],$$

and the map $E_1^{11} \to E_1^{21}$ can be identified with the differential operator $f \mapsto df$. Since the characteristic of K is 0, $\mathrm{Ker}(E_1^{11} \to E_1^{21}) \cong \langle e(C)\rangle_K$. On the other hand,

$$E_1^{01} = H^0(S^o, R^1f_*\mathscr{O}_{U^o}) \longrightarrow E_1^{11} = H^0(S^o, R^1f_*\Omega^1_{U^o/S^o})$$

is zero because the Hodge-de Rham spectral sequence

$$E_1^{pq} = R^qf_*\Omega^p_{X/S} \Longrightarrow R^{p+q}f_*\Omega^\bullet_{X/S}$$

degenerates at E_1-terms. Thus we have $E_2^{11} = \mathrm{Ker}(E_1^{11} \to E_1^{21}) \cong \langle e(C)\rangle_K$. This completes the proof of (3).

Next we show (4). We consider the spectral sequence

$$E_1^{pq} = H^q(\overline{U}, \Omega^p_{\overline{U}}) \Longrightarrow H^{p+q}_{\mathrm{dR}}(\overline{U}/K).$$

Since $\overline{S}$ is affine, we have

$$V_{\mathrm{dR}} = E_3^{20} = E_\infty^{20}, \quad E_2^{11} = E_\infty^{11}, \quad E_2^{02} = 0$$

by the same argument as before. Hence it is enough to show E_2^{11} is generated by the cycle classes of $e(C)$ and $\widetilde{D}$ (as K-module). Let $j : U^o \hookrightarrow \overline{U}$ be the open immersion. Consider a commutative diagram

$$\begin{tikzcd}
\Gamma(\overline{U}, j_*\Omega^1_{U^o}/\Omega^1_{\overline{U}}) \arrow[r, "\delta"] & H^1(\overline{U}, \Omega^1_{\overline{U}}) \arrow[r, "j^*"] \arrow[d, "d"] & H^1(U^o, \Omega^1_{U^o}) \arrow[d, "d^o"] \\
 & H^1(\overline{U}, \Omega^2_{\overline{U}}) \arrow[r] & H^1(U^o, \Omega^2_{U^o})
\end{tikzcd} \qquad \text{(exact)}$$

Let $x \in \mathrm{Ker} d$. Then $j^*(x) \in \mathrm{Ker} d^o$. As we have seen in the proof of (3), the kernel of d^o is generated by the cycle class $[e(C)]$ as K-module. Therefore $x' := x - c[e(C)]$ for some $c \in K$ is contained in $\mathrm{Ker}(j^*) = \mathrm{Im}\ \delta$. However, as is well-known, the image of δ is generated by the cycle classes of the irreducible components of $\overline{U} - U^o$. This shows that x is a linear combination of the cycle classes of $e(C)$ and D. Since $\mathrm{Ker}(d)$ is generated by the cycle classes of $e(C)$ and D as K-module, so is E_2^{11}.

Next we show (1) and (5). Let $D = \sum_i D_i$ be the irreducible decomposition (note we assumed $K = \overline{K}$ in this proof). The composition

$$\langle e(C), \widetilde{D} \rangle_K \longrightarrow H^2_{\mathrm{dR}}(\overline{U}/K) \longrightarrow \bigoplus_i H^2_{\mathrm{dR}}(D_i) \qquad (3.2)$$

is given by intersection pairing on $\langle e(C), \widetilde{D} \rangle_K$. Then it is not hard to show that the kernel of (3.2) is one dimensional generated by the cycle class of $f^{-1}(t)$ for $t \in T_1$. Conversely $f^{-1}(t)$ goes to zero via the map $\langle e(C), \widetilde{D} \rangle_K \to H^2_{\mathrm{dR}}(\overline{U}/K)$. Summarizing this, we have a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & V_{\mathrm{dR}} \arrow[r] & H^2_{\mathrm{dR}}(\overline{U}/K) \arrow[r] & E_\infty^{11} \arrow[r] \arrow[d] & 0 \\
 & & \mathrm{Im}\langle e(C), \widetilde{D} \rangle_K \arrow[u, "\cup"] \arrow[ur, "\cong"] \arrow[r, "\subset"] & \bigoplus_i H^2_{\mathrm{dR}}(D_i) &
\end{tikzcd} \qquad \text{(exact)} \qquad (3.3)$$

with an exact row. This shows (5) because $\bigoplus_i H^2_{\mathrm{dR}}(D_i) = H^2_{\mathrm{dR}}(D)$. Let us show (1). We first note that the image of $\mathrm{NF}_{\mathrm{dR}}(X)^\perp$ via the natural map $H^2_{\mathrm{dR}}(X/K) \to H^2_{\mathrm{dR}}(\overline{U}/K)$ is contained in V_{dR}. In fact, this follows from (3.3) and the fact that the composition

$$\mathrm{NF}_{\mathrm{dR}}(X)^\perp \longrightarrow H^2_{\mathrm{dR}}(X/K) \longrightarrow H^2_{\mathrm{dR}}(\overline{U}/K) \longrightarrow \bigoplus_i H^2_{\mathrm{dR}}(D_i)$$

is zero. Now we have a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & V_{\mathrm{dR}} \arrow[r] & H^2_{\mathrm{dR}}(\overline{U}/K) \arrow[r] & \bigoplus_i H^2_{\mathrm{dR}}(D_i) & \\
0 \arrow[r] & \mathrm{NF}_{\mathrm{dR}}(X)^{\perp} \arrow[r] \arrow[u] & H^2_{\mathrm{dR}}(X/K) \arrow[r] \arrow[u, "a"] & \mathrm{NF}_{\mathrm{dR}}(X) \arrow[r] \arrow[u, "b"] & 0
\end{tikzcd} \tag{3.4}$$

with exact rows. Since $X - \overline{U} = E$ are additive fibers, a is surjective. Therefore it is enough to show that $\mathrm{Ker}(a) \to \mathrm{Ker}(b)$ is bijective. $\mathrm{Ker}(a)$ is the sub K-module generated by the irreducible components of E. This implies $\mathrm{NF}_{\mathrm{dR}}(X)^{\perp} \cap \mathrm{Ker}(a) = 0$ and hence $\mathrm{Ker}(a) \to \mathrm{Ker}(b)$ is injective. On the other hand, since $\mathrm{Ker}(b)$ is generated by the irreducible components of E, $\mathrm{Ker}(a) \to \mathrm{Ker}(b)$ is surjective. This completes the proof of (1).

Finally we prove (2). Let us consider a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & V_{\mathrm{PF}} \arrow[r] & H^2_{\mathrm{dR}}(U^o/K) \arrow[r] & \langle e(C) \rangle_K \arrow[r] & 0 \\
0 \arrow[r] & V_{\mathrm{dR}} \arrow[r] \arrow[u] & H^2_{\mathrm{dR}}(\overline{U}/K) \arrow[r] \arrow[u, "a"] & \mathrm{Im}\langle e(C), \widetilde{D} \rangle_K \arrow[r] \arrow[u, "b"] & 0
\end{tikzcd}$$

with exact rows. $\mathrm{Ker}(a)$ is the sub K-module generated by the irreducible components of D. Therefore $\mathrm{Ker}(a) \to \mathrm{Ker}(b)$ is surjective. Moreover $V_{\mathrm{dR}} \cap \mathrm{Ker}(a) = 0$ because $\mathrm{Ker}(a)$ has an injection into $\oplus_i H^2_{\mathrm{dR}}(D_i)$. Therefore $\mathrm{Ker}(a) \to \mathrm{Ker}(b)$ is injective and hence bijective. Clearly the map b is surjective, Since $\mathrm{Coker}(a) = H_1^{\mathrm{dR}}(D/K) \oplus H_1^{\mathrm{dR}}(f^{-1}(F)/K)$, (2) follows. $\square$

The following is straightforward from Prop. 3.4.

Proposition 3.5 *Suppose $K = \mathbb{C}$. Let $\mathrm{NF}_B(X) \subset H^2_B(X, \mathbb{Q}) = H^2_B(X(\mathbb{C}), \mathbb{Q})$ be the image of $\mathrm{NF}(X) \otimes \mathbb{Q}$ and $\mathrm{NF}_B(X)^{\perp}$ its orthogonal complement with respect to the cup-product pairing. Let $F^{\bullet}V_{\mathrm{PF}}$ denotes the Hodge filtration as in Thm.6.6 and $F^{\bullet}V_{\mathrm{dR}}$ the induced filtration via Prop.3.4 (2). Then there is the natural comparison isomorphism $\mathrm{NF}_B(X)^{\perp} \otimes \mathbb{C} \cong V_{\mathrm{dR}}$ and $\mathrm{NF}(X)^{\perp} := (\mathrm{NF}_B(X)^{\perp}, F^{\bullet}V_{\mathrm{dR}})$ forms the sub Hodge structure of $H^2(X, \mathbb{Q}) := (H^2_B(X(\mathbb{C}), \mathbb{Q}), F^{\bullet}H^2_{\mathrm{dR}}(X/\mathbb{C}))$.*

4 Explicit computations of regulator on K_1 of elliptic surfaces

We keep the notations in §3.1. We assume $K = \mathbb{C}$ and $E \neq \emptyset$ (cf. Prop. 3.4) in this section. $H_{\bullet}(X, \mathbb{Q}) = H_{\bullet}(X(\mathbb{C}), \mathbb{Q})$ denotes the Betti homology.

4.1 1-Extension of MHS's arising from a multiplicative fiber

For each $\gamma \in H_1(D, \mathbb{Q})$, there is a corresponding element $\xi_\gamma \in H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$ which is unique up to the decomposable part. It is given in the following way. Let $D_k = \sum_{i=1}^{n} D_k^{(i)} =$

$f^{-1}(P_k)$ be a multiplicative fiber over a point $P_k \in C$ and Q_i the intersection points. There are rational functions f_i on $D_k^{(i)}$ such that $\mathrm{Div}_{D_k^{(i)}}(f_i) = Q_{i+1} - Q_i$. Then we put

$$\xi_{D_k} := \sum_{i=1}^{n} [f_i, D_k^{(i)}] \in H^3_{\mathscr{M}}(X, \mathbb{Q}(2)).$$

If $\gamma = (m_1, \cdots, m_s) \in H_1(D, \mathbb{Z}) = \bigoplus_k H_1(D_k, \mathbb{Z}) \cong \mathbb{Z}^{\oplus k}$ then we put $\xi_\gamma := m_1\xi_{D_1} + \cdots + m_s\xi_{D_s}$. This depends on the choice of f_i's, though the ambiguity is killed by the decomposable part.



Let us recall the regulator $\mathrm{reg}(\xi_\gamma)$ from §2.2. Let

$$\mathrm{NF}^B(X) := \mathrm{Im}(H_2(D, \mathbb{Q}) \oplus H_2(E, \mathbb{Q}) \oplus H_2(e(C), \mathbb{Q}) \longrightarrow H_2(X, \mathbb{Q})).$$

The exact sequence

$$0 \longrightarrow H_2(X, \mathbb{Q})/H_2(D, \mathbb{Q}) \longrightarrow H_2(X, D, \mathbb{Q}) \longrightarrow H_1(D, \mathbb{Q}) \longrightarrow 0 \tag{4.1}$$

of mixed Hodge structures gives rise to a map

$$\begin{aligned} \rho : H_1(D, \mathbb{Q}) &\longrightarrow \mathrm{Ext}^1(\mathbb{Q}, H_2(X, \mathbb{Q})/H_2(D, \mathbb{Q})) \\ &\longrightarrow \mathrm{Ext}^1(\mathbb{Q}, H_2(X, \mathbb{Q})/\mathrm{NF}^B(X)) \\ &\xleftarrow{\cong} \mathrm{Ext}^1(\mathbb{Q}, \mathrm{NF}(X)^\perp \otimes \mathbb{Q}(2)) \end{aligned}$$

where $\mathrm{NF}(X)^\perp \subset H^2(X, \mathbb{Q})$ is the Hodge structure as in Prop.3.5. Then we have from Thm.2.1

$$\mathrm{reg}(\xi_\gamma) = \pm\rho(\gamma). \tag{4.2}$$

In this section, we shall use a slightly different 1-extension of mixed Hodge structures from (4.1).

Lemma 4.1 *$H_2(\overline{U}, D; \mathbb{Q}) \to H_1(D, \mathbb{Q})$ is surjective.*

Proof. The assertion is equivalent to saying that $H^1(\overline{U}, D; \mathbb{Q}) \to H^1(\overline{U}, \mathbb{Q})$ is surjective. Since the functional j-invariant of U/S is not constant (by the assumption), one has

$$\varGamma(S, R^1 f_* \mathbb{Q}) = H^1(f^{-1}(t), \mathbb{Q})^{\pi_1(S,t)} = 0$$

and hence $H^1(U, \mathbb{Q}) = H^1(S, \mathbb{Q})$. This and a commutative diagram

$$\begin{tikzcd}
0 = H^1_D(\overline{U}, \mathbb{Q}) \arrow[r] & H^1(\overline{U}, \mathbb{Q}) \arrow[r] & H^1(U, \mathbb{Q}) \arrow[r] & H^2_D(\overline{U}, \mathbb{Q}) \cong H_2(D, \mathbb{Q}) \\
0 = H^1_{T_1}(\overline{S}, \mathbb{Q}) \arrow[r] \arrow[u] & H^1(\overline{S}, \mathbb{Q}) \arrow[r] \arrow[u] & H^1(S, \mathbb{Q}) \arrow[r] \arrow[u, "\cong"] & H^2_{T_1}(\overline{S}, \mathbb{Q}) \cong H_0(T_1, \mathbb{Q}) \arrow[u, "\cup"]
\end{tikzcd}$$

yield

$$H^1(\overline{S}, \mathbb{Q}) \xrightarrow{\cong} H^1(\overline{U}, \mathbb{Q}). \tag{4.3}$$

Therefore, to show the surjectivity of $H^1(\overline{U}, D; \mathbb{Q}) \to H^1(\overline{U}, \mathbb{Q})$ it is enough to show that $H^1(\overline{S}, T_1; \mathbb{Q}) \to H^1(\overline{S}, \mathbb{Q})$ is surjective. However it is clear because $H^1(T_1, \mathbb{Q}) = 0$. $\square$

Using the isomorphisms

$$H_2(\overline{U}, \mathbb{Q})/H_2(D) \xrightarrow{\cong} H_2(X, \mathbb{Q})/\mathrm{NF}^B(X) \xleftarrow{\cong} \mathrm{NF}(X)^\perp \otimes \mathbb{Q}(2) \tag{4.4}$$

we have a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & H_2(X, \mathbb{Q})/H_2(D, \mathbb{Q}) \arrow[r] \arrow[d, "\text{surj.}"'] \arrow[dr, phantom, "\square"] & H_2(X, D; \mathbb{Q}) \arrow[r] \arrow[d, "\text{surj.}"'] & H_1(D, \mathbb{Q}) \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & H_2(X, \mathbb{Q})/\mathrm{NF}^B(X) \arrow[r] & \overline{H_2(X, D; \mathbb{Q})} \arrow[r] & H_1(D, \mathbb{Q}) \arrow[r] & 0 \\
0 \arrow[r] & H_2(\overline{U}, \mathbb{Q})/H_2(D) \arrow[r] \arrow[u, "\cong"] \arrow[d, equal, "\wr"'] & H_2(\overline{U}, D; \mathbb{Q}) \arrow[r] \arrow[u, "\cong"] & H_1(D, \mathbb{Q}) \arrow[r] \arrow[u, equal] & 0 \\
& \mathrm{NF}(X)^\perp \otimes \mathbb{Q}(2) & & &
\end{tikzcd}$$

Let

$$\Phi : H_2(\overline{U}, \mathbb{Z}) \longrightarrow \mathrm{Hom}(H^2_{\mathrm{dR}}(\overline{U}/\mathbb{C}), \mathbb{C}), \quad \Delta \longmapsto \left[\omega \mapsto \int_\Delta \omega \right]$$

be the period map. It follows from Prop.3.4 that Φ induces the following commutative diagram.

$$\begin{tikzcd}
H_2(\overline{U}, \mathbb{C}) \arrow[r, "\cong"] \arrow[d] & \mathrm{Hom}(H^2_{\mathrm{dR}}(\overline{U}/\mathbb{C}), \mathbb{C}) \arrow[d] \\
H_2(\overline{U}, \mathbb{C})/H_2(D) \arrow[r, "\cong"] & \mathrm{Hom}(V_{\mathrm{dR}}, \mathbb{C})
\end{tikzcd} \tag{4.5}$$

Hence we have

$$\mathrm{Ext}^1(\mathbb{Q}, \mathrm{NF}(X)^\perp \otimes \mathbb{Q}(2)) \cong \mathrm{Coker}\left[\Phi : H_2(\overline{U}, \mathbb{Q}) \to \mathrm{Hom}(F^1V_{\mathrm{dR}}, \mathbb{C})\right] \tag{4.6}$$

(cf. (2.16), (2.17)). Under this identification, we have

$$\mathrm{reg}(\xi_\gamma) = \pm\left[\omega \longmapsto \int_\Gamma \omega\right] \in \mathrm{Hom}(F^1V_{\mathrm{dR}}, \mathbb{C})/\mathrm{Im}\Phi \tag{4.7}$$

for $\Gamma \in H_2(\overline{U}, D, \mathbb{Q})$ satisfying $\gamma = \partial(\Gamma)$.

4.2 $\mathbb{E}(\overline{U}, D; \mathbb{Z})$ and $\mathbb{E}(\overline{U}, \mathbb{Z})$

Take a path $\gamma : [0,1] \to \overline{S}(\mathbb{C})$, $t \mapsto \gamma_t$ such that $\gamma_t \in S(\mathbb{C})$ for $t \neq 0, 1$. Take a cycle $\varepsilon \in H_1(f^{-1}(\gamma_{t_0}), \mathbb{Z})$ for some (fixed) $t_0 \in [0,1]$. Then it extends to a flat section $\varepsilon_t \in H_1(f^{-1}(\gamma_t), \mathbb{Z})$ over $t \in [0,1]$ in a unique way. We denote by $\Gamma(\varepsilon, \gamma)$ the fibration over the path γ whose fiber is ε_t.



Then

$$\Gamma(\varepsilon, \gamma) \in H_2(\overline{U}, f^{-1}(\gamma_0) \cup f^{-1}(\gamma_1); \mathbb{Z}),$$
$$\text{with} \quad \partial(\Gamma(\varepsilon, \gamma)) = \varepsilon_1 - \varepsilon_0 \in H_1(f^{-1}(\gamma_0) \cup f^{-1}(\gamma_1), \mathbb{Z}).$$

Define $\mathbb{E}(\overline{U}, D; \mathbb{Z}) \subset H_2(\overline{U}, D; \mathbb{Z})$ the subgroup generated by $\Gamma(\varepsilon, \gamma)$'s where γ and ε run over as above such that $\gamma_0, \gamma_1 \in T_1 = \overline{S} - S$. Define $\mathbb{E}(\overline{U}, \mathbb{Z})$ by an exact sequence

$$0 \longrightarrow \mathbb{E}(\overline{U}, \mathbb{Z}) \longrightarrow \mathbb{E}(\overline{U}, D; \mathbb{Z}) \longrightarrow H_1(D, \mathbb{Z}).$$

Proposition 4.2

$$\mathbb{E}(\overline{U}, D; \mathbb{Q}) \xrightarrow{\cong} H_2(\overline{U}, D; \mathbb{Q}), \tag{4.8}$$

$$\mathbb{E}(\overline{U}, \mathbb{Q}) \xrightarrow{\cong} H_2(\overline{U}, \mathbb{Q})/H_2(D, \mathbb{Q}) \cong \mathrm{NF}_B(X)^\perp \otimes \mathbb{Q} \tag{4.9}$$

where $\mathbb{E}(\overline{U},D;\mathbb{Q}) := \mathbb{E}(\overline{U},D;\mathbb{Z})\otimes\mathbb{Q}$ *etc. Hence we have*

$$\begin{tikzcd}
0 \arrow[r] & \mathbb{E}(\overline{U};\mathbb{Q}) \arrow[r] \arrow[d, "\cong"] & \mathbb{E}(\overline{U},D;\mathbb{Q}) \arrow[r] \arrow[d, "\cong"] & H_1(D,\mathbb{Q}) \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & H_2(\overline{U},\mathbb{Q})/H_2(D,\mathbb{Q}) \arrow[r] & H_2(\overline{U},D;\mathbb{Q}) \arrow[r] & H_1(D,\mathbb{Q}) \arrow[r] & 0.
\end{tikzcd}$$

Lemma 4.3 *The sequence*

$$H_2(U,\mathbb{Q}) \longrightarrow H_2(\overline{U},D;\mathbb{Q}) \longrightarrow H_1(D,\mathbb{Q}) \longrightarrow 0 \tag{4.10}$$

is exact.

Proof. We already showed the surjectivity of the right arrow (Lem. 4.1). It is enough to show

$$\mathrm{Im}(H_2(U,\mathbb{Q}) \longrightarrow H_2(\overline{U},D;\mathbb{Q})) = \mathrm{Im}(H_2(\overline{U},\mathbb{Q}) \longrightarrow H_2(\overline{U},D;\mathbb{Q})). \tag{4.11}$$

Consider a diagram

$$\begin{tikzcd}
 & H_2(D,\mathbb{Q}) \arrow[d, "a"] & & & \\
H_2(U,\mathbb{Q}) \arrow[r] & H_2(\overline{U},\mathbb{Q}) \arrow[r, "b"] \arrow[d] & H^2(D,\mathbb{Q}) \arrow[r] & H_1(U,\mathbb{Q}) \arrow[r, "c"] & H_1(\overline{U},\mathbb{Q}) \\
 & H_2(\overline{U},D;\mathbb{Q}) & & &
\end{tikzcd}$$

with exact row and column. Hence it is enough to show $\mathrm{Im}(ba) = \mathrm{Im}(b)$ or equivalently $\dim\mathrm{Coker}(ba) = \dim\mathrm{Coker}(b) (= \dim\mathrm{Ker}(c))$. Since ba is given by the intersection pairing, a direct calculation shows that $\dim\mathrm{Coker}(ba) = \dim H_0(T_1)$. On the other hand,

$$\mathrm{Ker}(H_1(U,\mathbb{Q}) \overset{c}{\to} H_1(\overline{U},\mathbb{Q})) \cong \mathrm{Ker}(H_1(S,\mathbb{Q}) \to H_1(\overline{S},\mathbb{Q})) \cong H_0(T_1)$$

(cf. (4.3)), and hence the assertion follows. □

Proof of Prop.4.2. Let $\mathscr{L}$ be the local system on $S(\mathbb{C})$ whose fiber is $H_1(f^{-1}(t),\mathbb{Q})$. Then the image of $H_2(U,\mathbb{Q})$ in $H_2(\overline{U},D;\mathbb{Q})$ coincides with that of $H_1(S,\mathscr{L})$. The homology group $H_1(S,\mathscr{L})$ is generated by $\Gamma(\varepsilon,\gamma)$'s such that $\gamma_0 = \gamma_1 \in S$ and $\varepsilon_0 = \varepsilon_1$. Take a path δ such that $\delta_0 \in T_1 = \overline{S} - S$ and $\delta_1 = \gamma_0 = \gamma_1$. Put $\widetilde{\gamma} = \delta\cdot\gamma\cdot\delta^{-1}$. Then the image of $\Gamma(\varepsilon,\gamma)$ in $H_2(\overline{U},D;\mathbb{Q})$ coincides with $\Gamma(\varepsilon,\widetilde{\gamma})$, and this is an element of $\mathbb{E}(\overline{U},D;\mathbb{Q})$. There remains to show the surjectivity of $\mathbb{E}(\overline{U},D;\mathbb{Q}) \to H_1(D,\mathbb{Q})$ (this gives an alternative proof of Lem.4.1). To do this, it is enough to show that for each $p \in T_1$, there is a path ν such that $\nu_0 = p$ and $\nu_1 \in T_1$, and there is a cycle $\alpha_t \in H_1(f^{-1}(\nu_t),\mathbb{Q})$ such that $\alpha_0 \neq 0$ and $\alpha_1 = 0$. Since

$$\partial(\Gamma(\alpha,\nu)) = (\cdots,0,\alpha_0,0,\cdots) \in H_1(D,\mathbb{Q}) = \bigoplus_{x\in T_1} H_1(f^{-1}(x),\mathbb{Q})$$

this implies the surjectivity of $\mathbb{E}(\overline{U}, D; \mathbb{Q}) \to H_1(D, \mathbb{Q})$. Fix paths ν' and ν'' such that $\nu'_0 = p$, $\nu'_1 = \nu''_1 = q \in S$ and $\nu''_0 \in T_1$. Fix $\alpha' \in H_1(f^{-1}(q), \mathbb{Q})$ such that α' goes to a nonzero cycle as $q \to p$, and $\alpha'' \in H_1(f^{-1}(q), \mathbb{Q})$ such that α'' goes to zero as $q \to \nu''_0$. Since $H_1(f^{-1}(q), \mathbb{Q})$ is an irreducible $\mathbb{Q}[\pi_1(S, q)]$-module, there is $g \in \mathbb{Q}[\pi_1(S, q)]$ such that $g(\alpha') = \alpha''$. Joining $\Gamma(\alpha', \nu')$, $\Gamma(\alpha', g)$ and $\Gamma(\alpha'', \nu'')$, we obtain $\Gamma(\alpha, \nu)$ as desired. This completes the proof of Prop.4.2.

4.3 Explicit computation of regulator on $H^3_{\mathscr{M}}(X, \mathbb{Q}(2))$

We summarize all of the results in §4.1 and §4.2 together with Thm.2.1 in the following theorem.

Theorem 4.4 *Let us keep the notations as in §3.1 and 4.2. Suppose $K = \mathbb{C}$ and $E \neq \emptyset$*[1]*. Then we have*

$$\mathrm{Ext}^1(\mathbb{Q}, \mathrm{NF}(X)^{\perp} \otimes \mathbb{Q}(2)) \cong \mathrm{Coker}\left[\Phi : \mathbb{E}(\overline{U}, \mathbb{Q}) \longrightarrow \mathrm{Hom}(F^1 V_{\mathrm{dR}}, \mathbb{C})\right]$$

where

$$\Phi(\Delta) = \left[\omega \longmapsto \int_{\Delta} \omega\right], \quad \omega \in F^1 V_{\mathrm{dR}}.$$

For $\gamma \in H_1(D, \mathbb{Q})$, choose $\Gamma \in \mathbb{E}(\overline{U}, D; \mathbb{Q})$ with $\partial(\Gamma) = \gamma$. Then

$$\mathrm{reg}(\gamma) = \pm \left[\omega \longmapsto \int_{\Gamma} \omega\right] \in \mathrm{Hom}(F^1 V_{\mathrm{dR}}, \mathbb{C})/\mathrm{Im}\Phi.$$

The point is that "$\omega \in F^1 V_{\mathrm{dR}}$" is an algebraic differential 2-form, and one can obtain a basis of V_{dR} and $F^1 V_{\mathrm{dR}}$ explicitly. Indeed, we embed V_{dR} into V_{PF} the inhomogeneous de Rham cohomology (cf. Prop. 3.3). Since the Picard-Fuchs operator can be written down explicitly (Cor. 6.2), one can obtain the basis in a down-to-earth manner (see also §6.3). We also need to compute a basis of $\mathbb{E}(\overline{U}, D; \mathbb{Q})$. To do this, we first choose a basis of $\mathbb{E}(\overline{U}, D; \mathbb{Q})/\mathbb{E}(\overline{U}, \mathbb{Q})$ by computing the boundary map $\partial : \mathbb{E}(\overline{U}, D; \mathbb{Q}) \to H_1(D, \mathbb{Q})$. To obtain a basis of $\mathbb{E}(\overline{U}, \mathbb{Q})$ we assume that the precision of the values of integrations can be raised as many as one likes. Then, by using the basis of V_{dR} together with (4.5) and Prop. 4.2, one can prove the linear independence of given cycles in $\mathbb{E}(\overline{U}, \mathbb{Q})$ if they were linear independent. Hence one can eventually obtain a basis of $\mathbb{E}(\overline{U}, \mathbb{Q})$.

We shall apply the above method to an example in the next section.

5 Example : $3y^2 + x^3 + (3x + 4t^l)^2$

Let $l \geq 1$ be an integer. We consider a minimal elliptic surface

$$f : X \longrightarrow \mathbb{P}^1, \quad f^{-1}(t) : 3y^2 + x^3 + (3x + 4t^l)^2 = 0$$

---

[1] This seems not to be a serious condition.

defined over $\mathbb{Q}$. There is the section $e : \mathbb{P}^1 \to X$ of "∞". Write $X_{\mathbb{C}} := X \times_{\mathbb{Q}} \mathbb{C}$.

The purpose of this section is to compute the real regulator

$$\mathrm{reg}_{\mathbb{R}} : H^3_{\mathscr{M}}(X, \mathbb{Q}(2)) \longrightarrow \mathrm{Ext}^1_{\mathbb{R}\text{-MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty = 1},$$

where $H^2(X)_{\mathrm{ind}} := H^2(X_{\mathbb{C}}, \mathbb{Q}(1))/\mathrm{NS}(X_{\mathbb{C}})$, especially for an element

$$\xi_{D_1} = \left[\frac{y-(x+4)}{y+(x+4)}, D_1\right] \in H^3_{\mathscr{M}}(X, \mathbb{Q}(2)) \quad (D_1 := f^{-1}(1))$$

arising from a split multiplicative fiber D_1 of type I_1. We note that if $(l, 6) = 1$ then ξ_{D_1} is an "integral" element, in the sense that it comes from the motivic cohomology of a proper flat regular model of X over $\mathbb{Z}$ (see [Sch] 1.1.6 for "unconditional" definition of integral elements).

5.1 Basic data of X

The following is easy to show (the proof is left to the reader).

- The Hodge numbers are as follows:

$$h^{10} = h^{01} = 0, \quad h^{20} = h^{02} = \lfloor \frac{l-1}{3} \rfloor, \quad h^{11} = 10(1 + h^{20}).$$

  In particular, $H^2_B(X)_{\mathrm{ind}} := H^2_B(X(\mathbb{C}), \mathbb{Q})/\mathrm{NS}(X_{\mathbb{C}}) \otimes \mathbb{Q} \neq 0$ if and only if $l \geq 4$. (If $1 \leq l \leq 3$, then X is a rational surface.)

- There are $(l+1)$-multiplicative fibers:

$$D_0 := f^{-1}(0)(\text{=type } \mathrm{I}_{3l}), \quad D_i := f^{-1}(\zeta_l^{i-1})(\text{=type } \mathrm{I}_1),$$

  where $\zeta_l = \exp(2\pi i/l)$ and $1 \leq i \leq l$. Moreover $D_1 = f^{-1}(1)$ is the unique split multiplicative fiber.

- If $3|l$, then there is no additive fiber. If $(3, l) = 1$ then $E = f^{-1}(\infty)$ is the unique additive fiber (type IV* if $l \equiv 1$ mod 3, and type IV if $l \equiv 2$ mod 3). In particular, $E \neq \emptyset$ if and only if $(l, 3) = 1$.

- $\mathrm{NF}(X_{\mathbb{C}}) \otimes \mathbb{Q} = \mathrm{NS}(X_{\mathbb{C}}) \otimes \mathbb{Q}$ if and only if l is odd ([St] Example 4).

Hereafter we assume $(l, 6) = 1$. Then

$$\mathrm{NF}_B(X_{\mathbb{C}})^{\perp} = \mathrm{NS}(X_{\mathbb{C}})^{\perp} \xrightarrow{\cong} H^2_B(X)_{\mathrm{ind}} := H^2_B(X(\mathbb{C}), \mathbb{Q}(1))/\mathrm{NS}(X_{\mathbb{C}}) \otimes \mathbb{Q}.$$

Moreover one has

$$\dim \mathbb{E}(\overline{U}, \mathbb{Q}) \overset{(4.9)}{=} \dim \mathrm{NF}(X)^{\perp} = l - 1, \tag{5.1}$$

$$\dim \mathbb{E}(\overline{U}, D; \mathbb{Q}) = (l-1) + \dim H_1(D, \mathbb{Q}) = 2l - 1. \tag{5.2}$$

5.2 Picard-Fuchs operator and V_{dR}

Put $D = D_0 + \cdots + D_l$, $U = X - (D + E)$, $\overline{U} = X - E$ and

$$V_{\mathrm{dR}} = \mathrm{Im}(\mathit{\Gamma}(\overline{U}, \Omega^2_{\overline{U}}) \to H^2_{\mathrm{dR}}(\overline{U}/K)), \quad V_{\mathrm{PF}} = \mathrm{Im}(\mathit{\Gamma}(U, \Omega^2_U) \to H^2_{\mathrm{dR}}(U/K))$$

as in §3.1 (in this case $U = U^o$). The Picard-Fuchs operator is

$$PF^* = \frac{6}{l}t(t^l - 1)\frac{d^2}{dt^2} + \frac{6}{l}((l+1)t^l - 1)\frac{d}{dt} + \frac{4l}{3}t^{l-1}$$

and letting $A = \mathbb{Q}[t, 1/(t(t^l - 1))]$, we have a canonical isomorphism

$$V_{\mathrm{PF}} \cong \mathrm{Coker}(A \xrightarrow{PF} Adt\frac{dx}{y}), \quad \text{with } PF : f \mapsto PF^*(f)dt\frac{dx}{y}.$$

The subspace $V_{\mathrm{dR}} \subset V_{\mathrm{PF}}$ and the Hodge filtration $F^\bullet V_{\mathrm{dR}}$ are given as follows (cf. Example 6.7):

$$V_{\mathrm{dR}} = \langle t^{i-1}dt\frac{dx}{y} \mid 1 \le i \le l - 1\rangle_{\mathbb{Q}} \tag{5.3}$$

$$F^1V_{\mathrm{dR}} = \langle t^{i-1}dt\frac{dx}{y} \mid 1 \le i \le l - \lfloor\frac{l-1}{3}\rfloor - 1\rangle_{\mathbb{Q}} \tag{5.4}$$

$$F^2V_{\mathrm{dR}} = \langle t^{i-1}dt\frac{dx}{y} \mid 1 \le i \le \lfloor\frac{l-1}{3}\rfloor\rangle_{\mathbb{Q}} \tag{5.5}$$

5.3 Cycles Δ and Γ

Let δ_0 (resp. δ_1) be the homology cycle in $H_1(f^{-1}(t), \mathbb{Z})$ which vanishes as $t \to 0$ (resp. $t \to 1$). Define Δ and Γ to be fibrations over the segment $[0, 1] \subset \mathbb{P}^1(\mathbb{C})$ whose fibers are the vanishing cycles δ_1 and δ_0 respectively.

$$\Delta \in H_2(\overline{U}, D_0; \mathbb{Z}), \quad \Gamma \in H_2(\overline{U}, D_1; \mathbb{Z}).$$

The boundary $\partial\Delta$ (resp. $\partial\Gamma$) is a generator of the homology group $H_1(D_0, \mathbb{Z}) \cong \mathbb{Z}$ (resp. $H_1(D_1, \mathbb{Z}) \cong \mathbb{Z}$).

Figure of Δ



Figure of Γ



Let $r_1(t) < r_2(t) < r_3(t)$ be the real roots of $x^3 + (3x + 4t^l)^2$ for $0 < t < 1$. Then one has

$$\int_\Delta t^{j-1}dt\frac{dx}{y} = 2\sqrt{-3}\overbrace{\int_0^1 t^{j-1}dt\int_{r_1(t)}^{r_2(t)}\frac{dx}{\sqrt{x^3+(3x+4t^l)^2}}}^{\text{positive real number}} \in \sqrt{-1}\mathbb{R}_{>0} \tag{5.6}$$

$$\int_\Gamma t^{j-1}dt\frac{dx}{y} = 2\sqrt{-3}\overbrace{\int_0^1 t^{j-1}dt\int_{r_2(t)}^{r_3(t)}\frac{dx}{\sqrt{x^3+(3x+4t^l)^2}}}^{\text{purely imaginary number}\neq 0} \in \mathbb{R}_{>0}. \tag{5.7}$$

Let $\sigma : X_{\mathbb{C}} \to X_{\mathbb{C}}$ be an automorphism given by $\sigma(x, y, t) = (x, y, \zeta_l t)$. Since

$$\sigma^* t^{j-1}dt\frac{dx}{y} = \zeta_l^j t^{j-1}dt\frac{dx}{y},$$

one has

$$\int_{\sigma_*^k\Delta} t^{j-1}dt\frac{dx}{y} = \zeta_l^{kj}\int_\Delta t^{j-1}dt\frac{dx}{y}.$$

This and an elementary calculation show that

$$\Delta - \sigma_*\Delta, \Delta - \sigma_*^2\Delta, \cdots, \Delta - \sigma_*^{l-1}\Delta$$

are linearly independent in $\mathbb{E}(\overline{U}, \mathbb{Q})$. Since it is $(l-1)$-dimensional by (5.1), the above is a basis of $\mathbb{E}(\overline{U}, \mathbb{Q})$. $\mathbb{E}(\overline{U}, D; \mathbb{Q})/\mathbb{E}(\overline{U}, \mathbb{Q})$ is l-dimensional with a basis

$$\Gamma, \sigma_*\Gamma, \cdots, \sigma_*^{l-1}\Gamma.$$

Let F_∞ denotes the infinite Frobenius morphism. Then

$$F_\infty(\Delta) = -\Delta, \quad F_\infty(\Gamma) = \Gamma.$$

Thm.4.4 together with the above results yields the following.

Theorem 5.1 *Suppose* $(l, 6) = 1$. *Put* $h = \dim F^1V_{\mathrm{dR}} = l - \lfloor \frac{l-1}{3} \rfloor - 1$ *and* $\zeta = \exp(2\pi i/l)$. *Let*

$$A = \left((\zeta^{pq} - \zeta^{-pq}) \int_\Delta t^{p-1}dt\frac{dx}{y} \right)_{1 \le p \le h,\ 1 \le q \le (l-1)/2}$$

be $h \times (l-1)/2$*-matrix (the entries are real numbers by* (5.6)*). Then*

$$\mathrm{Ext}^1_{\mathbb{R}\text{-MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty = 1} \cong \mathrm{Coker}[A : \mathbb{R}^{(l-1)/2} \longrightarrow \mathbb{R}^h].$$

and we have

$$\overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1}) = \pm \left(\int_\Gamma dt\frac{dx}{y}, \cdots, \int_\Gamma t^{h-1}dt\frac{dx}{y} \right) \in \mathbb{R}^h/\mathrm{Im}A$$

under the above isomorphism.

Corollary 5.2 *Suppose* $(l, 6) = 1$. *Then*

$$\overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1}) \neq 0 \in \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty = 1}$$

and hence ξ_{D_1} *is regulator indecomposable.*

Proof. Put $h = l - \lfloor \frac{l-1}{3} \rfloor - 1$ and $\zeta = \exp(2\pi i/l)$. Put

$$I_p := \int_\Delta t^{p-1}dt\frac{dx}{y}, \quad J_p := \int_\Gamma t^{p-1}dt\frac{dx}{y}.$$

Then

$$\overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1}) \neq 0 \in \mathrm{Ext}^1_{\mathrm{MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty = 1}$$

if and only if the rank of a matrix

$$\begin{pmatrix} (\zeta-\zeta^{-1})I_1 & (\zeta^2-\zeta^{-2})I_1 & \cdots & (\zeta^{(l-1)/2}-\zeta^{-(l-1)/2})I_1 & J_1 \\ (\zeta^2-\zeta^{-2})I_2 & (\zeta^4-\zeta^{-4})I_2 & \cdots & (\zeta^{l-1}-\zeta^{-(l-1)})I_2 & J_2 \\ \vdots & \vdots & & \vdots & \vdots \\ (\zeta^h-\zeta^{-h})I_h & (\zeta^{2h}-\zeta^{-2h})I_h & \cdots & (\zeta^{h(l-1)/2}-\zeta^{-h(l-1)/2})I_h & J_h \end{pmatrix} \tag{5.8}$$

is maximal. Thus it is enough to show that

$$\det\begin{pmatrix} (\zeta-\zeta^{-1}) & (\zeta^2-\zeta^{-2}) & \cdots & (\zeta^{(l-1)/2}-\zeta^{-(l-1)/2}) & J_1/I_1 \\ (\zeta^2-\zeta^{-2}) & (\zeta^4-\zeta^{-4}) & \cdots & (\zeta^{l-1}-\zeta^{-(l-1)}) & J_2/I_2 \\ \vdots & \vdots & & \vdots & \vdots \\ (\zeta^k-\zeta^{-k}) & (\zeta^{2k}-\zeta^{-2k}) & \cdots & (\zeta^{k(l-1)/2}-\zeta^{-k(l-1)/2}) & J_k/I_k \end{pmatrix} \neq 0 \tag{5.9}$$

where $k=(l+1)/2$. Since the sum of the $(k-1)$-th row and k-th row is $(0,\cdots,0,J_{k-1}/I_{k-1}+J_k/I_k)$, one has

$$\begin{aligned} \text{(5.9)} =& (J_{k-1}/I_{k-1}+J_k/I_k)\times\det(\zeta^{pq}-\zeta^{-pq})_{1\le p,q\le(l-1)/2} \\ =& (J_{k-1}/I_{k-1}+J_k/I_k)\times\sqrt{(-l)^{(l-1)/2}}. \end{aligned}$$

Since $J_p/I_p\in i\mathbb{R}_{>0}$ by (5.6) and (5.7), this is non-zero. □

5.4 Another description of $\int_\Delta$ and $\int_\Gamma$

When $l=1$, $f:X\to\mathbb{P}^1$ is the universal elliptic curve over $X_1(3)$. Using this, one can obtain another description of the real regulator.

Let $q=\exp(2\pi iz)$ and

$$E_{3,a}(z):=1-9\sum_{n=1}^{\infty}\left(\sum_{k|n}\left(\frac{k}{3}\right)k^2\right)q^n,$$

$$E_{3,b}(z):=\sum_{n=1}^{\infty}\left(\sum_{k|n}\left(\frac{n/k}{3}\right)k^2\right)q^n$$

be the Eisenstein series of weight 3 for $\Gamma_1(3)$, where $(\frac{k}{3})$ denotes the Legendre symbol. Then

$$t^l=\frac{E_{3,a}}{E_{3,a}+27E_{3,b}}$$

and

$$l\frac{dt}{t}\frac{dx}{y}=-27E_{3,b}\frac{du}{u}\frac{dq}{q},\qquad \frac{lt^{l-1}dt}{t^l-1}\frac{dx}{y}=E_{3,a}\frac{du}{u}\frac{dq}{q}$$

where “du/u” denotes the canonical invariant 1-form of the Tate curve around the cusp $z = i\infty$ ($t = 1$). Therefore we have

$$\int_\Delta t^{j-1}dt\frac{dx}{y} = \frac{-27}{l} \times (2\pi i)^2 \int_0^{i\infty} t^j E_{3b}(z)dz \tag{5.10}$$

$$\int_\Gamma t^{j-1}dt\frac{dx}{y} = \frac{-27}{l} \times (2\pi i)^2 \int_0^{i\infty} t^j E_{3b}(z)zdz. \tag{5.11}$$

On the other hand there are formulas

$$\frac{E_{3a}}{E_{3a}+27E_{3b}}(\frac{-1}{3z}) = \frac{27E_{3b}}{E_{3a}+27E_{3b}}(z), \quad 27E_{3b}(\frac{-1}{3z}) = 3\sqrt{3}iz^3E_{3a}(z) \tag{5.12}$$

on the Eisenstein series. Applying (5.12) to (5.10) and (5.11), we have the following theorem.

Theorem 5.3 *Put* $c := \exp(-2\pi/\sqrt{3}) = 0.026579933\cdots$. *Define rational numbers* $a_n(j)$ *and* $b_n(j)$ *by*

$$\begin{aligned} E_{3b}\left(\frac{E_{3a}}{E_{3a}+27E_{3b}}\right)^{j/l} &= \sum_{n=1}^\infty a_n(j)q^n \\ &= q + \left(3 - 27\frac{j}{l}\right)q^2 + \left(9 - \frac{81}{2}\frac{j}{l} + \frac{729}{2}\left(\frac{j}{l}\right)^2\right)q^3 + \cdots, \end{aligned}$$

$$\begin{aligned} E_{3a}\left(\frac{E_{3b}}{q(E_{3a}+27E_{3b})}\right)^{j/l} &= \sum_{n=0}^\infty b_n(j)q^n \\ &= 1 + \left(-9 - 15\frac{j}{l}\right)q + \left(27 + \frac{387}{2}\frac{j}{l} + \frac{225}{2}\left(\frac{j}{l}\right)^2\right)q^2 + \cdots. \end{aligned}$$

Put

$$\begin{aligned} I(j) &= \sum_{n=1}^\infty \frac{a_n(j)}{n}c^n + 3^{3j/l-3}\sum_{n=0}^\infty b_n(j)\left(\frac{1}{n+j/l} + \frac{\sqrt{3}}{2\pi(n+j/l)^2}\right)c^{n+j/l} \\ J(j) &= \sum_{n=1}^\infty a_n(j)\left(\frac{2\pi}{\sqrt{3}n} + \frac{1}{n^2}\right)c^n + 2\pi\cdot 3^{3j/l-7/2}\sum_{n=0}^\infty \frac{b_n(j)}{n+j/l}c^{n+j/l}. \end{aligned}$$

Then we have

$$\int_\Delta t^{j-1}dt\frac{dx}{y} = \frac{54\pi i}{l}I(j), \quad \int_\Gamma t^{j-1}dt\frac{dx}{y} = \frac{-27}{l}J(j)$$

for $1 \le j \le l-1$.

This is useful since the series $I(j)$ and $J(j)$ converge rapidly !

Example 5.4 Suppose $l = 5$. Then X is a K3 surface. By Thm.5.3, one has

	$I(j)$	$J(j)$
$j = 1$	0.42745977255318	0.214998778914541
$j = 2$	0.151180954233147	0.154831831238613
$j = 3$	0.0871841692346256	0.130967280361779
$j = 4$	0.0603840144077692	0.116576757544694

$$\mathrm{Ext}^1_{\mathbb{R}\text{-MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty = 1} \cong \mathrm{Coker}(\mathbb{R}^2 \xrightarrow{A} \mathbb{R}^3).$$

Since this is 1-dimensional, this has the canonical base $e_{\mathrm{ind},\mathbb{Q}}$ (up to $\mathbb{Q}^\times$) and a *different* base $e^{\mathit{false}}_{\mathrm{ind},\mathbb{Q}}$ (§2.4). With respect to $e^{\mathit{false}}_{\mathrm{ind},\mathbb{Q}}$, one has

$$\overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1}) = \pi^2 \begin{vmatrix} i(\zeta - \zeta^{-1})I(1) & i(\zeta^2 - \zeta^{-2})I(1) & J(1) \\ i(\zeta^2 - \zeta^{-2})I(2) & i(\zeta^4 - \zeta^{-4})I(2) & J(2) \\ i(\zeta^3 - \zeta^{-3})I(3) & i(\zeta^6 - \zeta^{-6})I(3) & J(3) \end{vmatrix} \mod \mathbb{Q}^\times \quad (\zeta := \exp(2\pi i/5)).$$

Since $s = (l-1)/2 = 2$ and $\det H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}} \otimes [\det H^2_B(X_{\mathbb{C}})_{\mathrm{ind}}]^{-1} = \sqrt{5}$ one has

$$\begin{aligned} \overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1}) &= \frac{\sqrt{5}}{\pi^2} \cdot \pi^2 \begin{vmatrix} i(\zeta - \zeta^{-1})I(1) & i(\zeta^2 - \zeta^{-2})I(1) & J(1) \\ i(\zeta^2 - \zeta^{-2})I(2) & i(\zeta^4 - \zeta^{-4})I(2) & J(2) \\ i(\zeta^3 - \zeta^{-3})I(3) & i(\zeta^6 - \zeta^{-6})I(3) & J(3) \end{vmatrix} \\ &= \pm 5\sqrt{5} I(1)I(2)I(3) \left(\frac{J(2)}{I(2)} + \frac{J(3)}{I(3)} \right) \\ &= 0.159139061341585 \mod \mathbb{Q}^\times \end{aligned}$$

with respect to $e_{\mathrm{ind},\mathbb{Q}}$ by Prop.2.2.

Example 5.5 Suppose $l = 7$. Then $h^{20}(X) = h^{02}(X) = 2$, $h^{11}(X) = 30$.

	$I(j)$	$J(j)$
$j = 1$	0.740059830730164	0.259822532138057
$j = 2$	0.24646699651114	0.179959563808791
$j = 3$	0.137265313181901	0.150366403148913
$j = 4$	0.0929578147374374	0.133576277272453
$j = 5$	0.0696363855176379	0.122069272279075
$j = 6$	0.0554349861351089	0.11332167027564

$$\mathrm{Ext}^1_{\mathbb{R}\text{-MHS}}(\mathbb{R}, H^2(X)_{\mathrm{ind}} \otimes \mathbb{R}(1))^{F_\infty = 1} \cong \mathrm{Coker}(\mathbb{R}^3 \xrightarrow{A} \mathbb{R}^4).$$

Since $s = (l-1)/2 = 3$ and $\det H^2_{\mathrm{dR}}(X/\mathbb{Q})_{\mathrm{ind}} \otimes [\det H^2_B(X_{\mathbb{C}})_{\mathrm{ind}}]^{-1} = \sqrt{-7}$, one has

$$\begin{aligned}\overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1}) &= \frac{\sqrt{7}}{\pi^3}\cdot\pi^3 \begin{vmatrix} i(\zeta-\zeta^{-1})I(1) & i(\zeta^2-\zeta^{-2})I(1) & i(\zeta^3-\zeta^{-3})I(1) & J(1) \\ i(\zeta^2-\zeta^{-2})I(2) & i(\zeta^4-\zeta^{-4})I(2) & i(\zeta^6-\zeta^{-6})I(2) & J(2) \\ i(\zeta^3-\zeta^{-3})I(3) & i(\zeta^6-\zeta^{-6})I(3) & i(\zeta^9-\zeta^{-9})I(3) & J(3) \\ i(\zeta^4-\zeta^{-4})I(4) & i(\zeta^8-\zeta^{-8})I(4) & i(\zeta^{12}-\zeta^{-12})I(4) & J(4) \end{vmatrix} \\ &= \pm 49 I(1)I(2)I(3)I(4)\left(\frac{J(3)}{I(3)}+\frac{J(4)}{I(4)}\right) \\ &= 0.288802269779652 \mod \mathbb{Q}^{\times} \quad (\zeta := \exp(2\pi i/7))\end{aligned}$$

with respect to the canonical $\mathbb{Q}$-structure $e_{\mathrm{ind},\mathbb{Q}}$.

Remark 5.6 *According to the Beilinson conjecture, $\overline{\mathrm{reg}}_{\mathbb{R}}(\xi_{D_1})$ in Example 5.4 or 5.5 is expected to be the value of the L-function $L(h^2(X)_{\mathrm{ind}}, s)$ at $s=1$* ([S]).

6 Appendix : Gauss-Manin connection and Picard-Fuchs operator

Let R be an integral domain of characteristic 0 in which 6 is invertible. For a smooth scheme Y over T, we denote by $\Omega^q_{Y/T} = \overset{q}{\wedge}_{\mathscr{O}_Y}\Omega^1_{Y/T}$ the sheaf of relative differential q-forms on Y over T. If $T = \mathrm{Spec} R$, we simply write $\Omega^q_Y = \Omega^q_{Y/R}$.

6.1 Explicit formulas

Let S be an irreducible affine smooth scheme over R of relative dimension one. Let $g_2, g_3 \in \mathscr{O}_S(S) = \Gamma(S, \mathscr{O}_S)$ satisfy $g_2^3 - 27g_3^2 \in \mathscr{O}_S(S)^{\times}$. Let $f : U \to S$ be a projective smooth family of elliptic curves whose affine form is given by a Weierstrass equation $y^2 = 4x^3 - g_2x - g_3$. More precisely letting

$$U_0 = \mathrm{Spec}\mathscr{O}_S(S)[x,y]/(y^2 - 4x^3 + g_2x + g_3),$$

$$U_\infty = \mathrm{Spec}\mathscr{O}_S(S)[u,z]/(z - 4u^3 + g_2uz^2 + g_3z^3),$$

U is obtained by gluing U_0 and U_∞ via identification $u = x/y,\ z = 1/y$. Let $e : S \longrightarrow U$ be a section given by $(u,z) = (0,0)$. We shall write down the *Gauss-Manin connection* (cf. [H] Ch.III, §4)

$$\nabla : H^1_{\mathrm{dR}}(U/S) \longrightarrow H^1_{\mathrm{dR}}(U/S) \otimes \Omega^1_S$$

(we use the same symbol "Ω^1_S" for $\Gamma(S, \Omega^1_S)$ since it will be clear from the context which is meant). To do this we fix a description of the Cech cohomology $H^q_{\mathrm{dR}}(U/S) := \check{H}^q(U, \Omega^{\bullet}_{U/S})$. We write

$$\check{C}^0(\mathscr{F}) := \Gamma(U_0, \mathscr{F}) \oplus \Gamma(U_\infty, \mathscr{F}), \quad \check{C}^1(\mathscr{F}) := \Gamma(U_0 \cap U_\infty, \mathscr{F})$$

for a (Zariski) sheaf $\mathscr{F}$. Then the double complex

$$\begin{tikzcd}
\check{C}^0(\mathscr{O}_U) \arrow[r, "d"] \arrow[d, "\delta"'] & \check{C}^0(\Omega^1_{U/S}) \arrow[d, "\delta"] \\
\check{C}^1(\mathscr{O}_U) \arrow[r, "d"] & \check{C}^1(\Omega^1_{U/S})
\end{tikzcd}
\qquad
\begin{tikzcd}
(x_0, x_\infty) \arrow[r, "d"] \arrow[d, "\delta"] & (dx_0, dx_\infty) \\
x_0 - x_\infty &
\end{tikzcd}$$

gives rise to the total complex

$$\check{C}^\bullet(U/S) : \check{C}^0(\mathscr{O}_U) \overset{\delta\oplus d}{\longrightarrow} \check{C}^1(\mathscr{O}_U) \oplus \check{C}^0(\Omega^1_{U/S}) \overset{(-d)\oplus\delta}{\longrightarrow} \check{C}^1(\Omega^1_{U/S})$$

of R-modules starting from degree 0, and the cohomology of it gives the de Rham cohomology $H^\bullet_{\mathrm{dR}}(U/S)$:

$$H^q_{\mathrm{dR}}(U/S) = H^q(\check{C}^\bullet(U/S)), \quad q \geq 0.$$

Elements of $H^1_{\mathrm{dR}}(U/S)$ are represented by cocycles

$$(f) \times (x_0, x_\infty) \quad \text{with } df = x_0 - x_\infty.$$

Theorem 6.1 *Suppose that Ω^1_S is a free $\mathscr{O}_S$-module with a base $dt \in \Gamma(S, \Omega^1_S)$. For $f \in \mathscr{O}_S(S)$, we define $f' \in \mathscr{O}_S(S)$ by $df = f'dt$. Let*

$$\omega := (0) \times (\frac{dx}{y}, \frac{dx}{y}) \tag{6.1}$$

$$\omega^* := (\frac{x^2}{y}) \times (\frac{xdx}{2y}, \frac{(2g_2x^2 + 3g_3x)dx}{2y^3}) \tag{6.2}$$

be elements in $H^1_{\mathrm{dR}}(U/S)$. Put $\Delta := g_2^3 - 27g_3^2$. Then we have

$$\nabla(\omega) = \left(\frac{-3(2g_2g_3' - 3g_3g_2')}{\Delta}\omega^* + \frac{\Delta'}{12\Delta}\omega \right) \otimes dt \in H^1_{\mathrm{dR}}(U/S) \otimes \Omega^1_S, \tag{6.3}$$

$$\nabla(\omega^*) = \left(-\frac{\Delta'}{12\Delta}\omega^* + \frac{g_2(2g_2g_3' - 3g_3g_2')}{16\Delta}\omega \right) \otimes dt \in H^1_{\mathrm{dR}}(U/S) \otimes \Omega^1_S. \tag{6.4}$$

Note that ω and ω^* are basis of the free $\mathscr{O}(S)$-module $H^1_{\mathrm{dR}}(U/S)$:

$$H^1_{\mathrm{dR}}(U/S) = \mathscr{O}_S(S)\omega \oplus \mathscr{O}_S(S)\omega^*.$$

Put $\mathscr{H} := R^1 f_* \Omega^\bullet_{U/S}$ and

$$\mathscr{H}^{1,0} := f_*\Omega^1_{U/S} = \mathscr{O}_S\omega, \quad \mathscr{H}^{0,1} := \mathscr{H}/\mathscr{H}^{1,0} \cong \mathscr{O}_S\omega^*.$$

The Gauss-Manin connection induces the following diagram

$$\begin{tikzcd}
 & 0 \arrow[d] \\
 & \mathscr{H}^{1,0} \otimes \Omega^1_S \arrow[d] \\
\mathscr{H} \arrow[r, "\nabla"] \arrow[d, "="'] & \mathscr{H} \otimes \Omega^1_S \arrow[d] \\
\mathscr{H} \arrow[r, "\widetilde{\nabla}"] & \mathscr{H}^{0,1} \otimes \Omega^1_S \arrow[d] \\
 & 0
\end{tikzcd} \tag{6.5}$$

Let

$$\overline{\nabla} : \mathscr{H}^{1,0} \longrightarrow \mathscr{H}^{0,1} \otimes \Omega^1_S \tag{6.6}$$

be the $\mathscr{O}_S$-linear map induced from the Gauss-Manin connection ∇. Let $S^o \subset S$ be a Zariski open set such that $\overline{\nabla}$ is bijective on S^o (cf. Prop.6.3). Then (6.5) gives rise to an exact sequence

$$\mathrm{Ker}\widetilde{\nabla}|_{S^o} \longrightarrow \mathscr{H}^{1,0}|_{S^o} \otimes \Omega^1_{S^o} \longrightarrow \mathrm{Coker}\nabla|_{S^o} \longrightarrow 0 \tag{6.7}$$

and the natural map $\mathrm{Ker}\widetilde{\nabla}|_{S^o} \to \mathscr{H}^{0,1}|_{S^o}$ is bijective. Hence we have an exact sequence

$$\begin{tikzcd}
0 \arrow[r] & \mathit{\Gamma}(S^o, \mathscr{H}^{0,1}) \arrow[r, "PF"] & \mathit{\Gamma}(S^o, \mathscr{H}^{1,0} \otimes \Omega^1_S) \arrow[r] & \mathit{\Gamma}(S^o, \mathrm{Coker}\nabla) \arrow[r] \arrow[d, equal] & 0 \\
 & & & H^1_{\mathrm{dR}}(S^o, \mathscr{H}). &
\end{tikzcd} \tag{6.8}$$

The map PF in (6.8) is called the *Picard-Fuchs operator*.

Corollary 6.2 *Suppose that $\Omega^1_{S^o}$ is a free $\mathscr{O}_{S^o}$-module with a base $dt \in \mathit{\Gamma}(S^o, \Omega^1_{S^o})$. Write $f'dt = df$ for $f \in \mathscr{O}(S^o)$. Put*

$$A := \frac{\Delta}{3(2g_2g_3' - 3g_2'g_3)}$$

$$B := \frac{1}{48}\left(\frac{-g_2(g_2')^2 + 12(g_3')^2}{2g_2g_3' - 3g_2'g_3} + 4\left(\frac{g_2^2g_2' - 18g_3g_3'}{2g_2g_3' - 3g_2'g_3}\right)'\right).$$

Then the Picard-Fuchs operator is described as follows.

$$PF(f\omega^*) = (f''A + f'(t)A' + fB)\omega \otimes dt, \quad f \in \mathscr{O}(S^o)$$

Cor. 6.2 follows from Thm. 6.1 by a direct (but rather long and tedious) calculation. The details are left to the reader.

The following is also straightforward from Thm. 6.1.

Proposition 6.3 *The $\mathscr{O}_S$-linear map* (6.6) *is described as follows.*

$$\overline{\nabla}(\omega) = \frac{-3(2g_2g_3' - 3g_3g_2')}{\Delta}\omega^* \otimes dt.$$

In particular, noting

$$\frac{j'}{j} = 27 \cdot \frac{g_3}{g_2} \cdot \frac{2g_2g_3' - 3g_2'g_3}{\Delta}, \quad j := \frac{1728g_2^3}{g_2^3 - 27g_3^2},$$

(6.6) *is bijective if and only if*

$$\frac{g_2}{g_3}\frac{dj}{j} \in \Omega_S^1$$

is a base of $\mathscr{O}_S$-module.

6.2 Proof of Theorem 6.1

Lemma 6.4

$$(\frac{x^i}{y^j}) \times (0,0) \equiv (0) \times (0, d(\frac{x^i}{y^j})) \quad (0 \leq i \leq j)$$

$$(\frac{x}{y}) \times (0,0) \equiv (0) \times (0, d(\frac{x}{y})) = (0) \times (0, (-\frac{1}{2} - \frac{g_2x}{y^2} - \frac{3g_3}{2y^2})\frac{dx}{y})$$

$$(\frac{1}{y}) \times (0,0) \equiv (0) \times (0, d(\frac{1}{y})) = (0) \times (0, \frac{-12x^2 + g_2}{2y^2}\frac{dx}{y})$$

$$(y) \times (0,0) \equiv (0) \times (-dy, 0) = (0) \times (\frac{g_2 - 12x^2}{2}\frac{dx}{y}, 0)$$

where "$\equiv$" denote modulo $\mathrm{Im}\check{C}^0(\mathscr{O}_U)$.

Proof. Straightforward from the definition. $\square$

Lemma 6.5 *Let η_U be the generic point of U. We think of dx and dt as elements in $\Gamma(\eta_U, \Omega_U^1)$. Then*

$$\frac{dx}{y} \in \Gamma(U_\infty, \Omega_U^1), \tag{6.9}$$

$$\frac{\widetilde{dx}}{y} := \frac{dx}{y} - \frac{(6g_2x^2 - 9g_3x - g_2^2)(g_2'x + g_3')}{\Delta}\frac{dt}{y} \in \Gamma(U_0, \Omega_U^1). \tag{6.10}$$

Proof.

$$\begin{aligned}\frac{dx}{y} &= du - \frac{u}{z}((12u^2 - g_2z^2)du - (2g_2uz + 3g_3z^2)dz - (g_2'uz^2 + g_3'z^3)dt) \\ &\equiv -\frac{12u^3}{z}du \mod \Gamma(U_\infty, \Omega_U^1) \\ &= -3(1 + g_2uz + g_3z^2)du \\ &\equiv 0 \mod \Gamma(U_\infty, \Omega_U^1).\end{aligned}$$

Hence (6.9) follows. Next we show (6.10). Since $f(x) = 4x^3 - g_2x - g_3$ is prime to $f'(x) = 12x^2 - g_2$, there are $a(x)$ and $b(x)$ such that

$$a(x)f(x) + b(x)f'(x) = 1.$$

Explicitly, they are given as follows.

$$a(x) = \frac{9(3g_3 - 2g_2x)}{\Delta}, \quad b(x) = \frac{6g_2x^2 - 9g_3x - g_2^2}{\Delta}.$$

Now

$$\begin{aligned}\frac{dx}{y} &= \frac{a(x)f(x)dx + b(x)f'(x)dx}{y} \\ &= \frac{a(x)y^2dx + b(x)(2ydy + (g_2'x + g_3')dt)}{y} \\ &= a(x)ydx + 2b(x)dy + \frac{b(x)(g_2'x + g_3')}{y}dt \in \mathit{\Gamma}(U_0 \cap U_\infty, \Omega_U^1)\end{aligned}$$

and hence we have

$$\frac{\widetilde{dx}}{y} = \frac{dx}{y} - \frac{b(x)(g_2'x + g_3')}{y}dt = a(x)ydx + 2b(x)dy \in \mathit{\Gamma}(U_0, \Omega_U^1).$$

□

Let us prove (6.3). Let

$$\widetilde{\omega} := (0) \times (\frac{\widetilde{dx}}{y}, \frac{dx}{y}) \in \check{C}^1(\mathscr{O}_U) \oplus \check{C}^0(\Omega_U^1)$$

be a lifting of ω where $\frac{\widetilde{dx}}{y}$ is as in (6.10). Let

$$\begin{tikzcd}
\check{C}^0(\mathscr{O}_U) \arrow[r, "d"] \arrow[d, "\delta"'] & \check{C}^0(\Omega_U^1) \arrow[r, "d"] \arrow[d, "\delta"] & \check{C}^0(\Omega_U^2) \arrow[d, "\delta"] \\
\check{C}^1(\mathscr{O}_U) \arrow[r, "d"] & \check{C}^1(\Omega_U^1) \arrow[r, "d"] & \check{C}^1(\Omega_U^2)
\end{tikzcd}$$

be the double complex and

$$\check{C}^0(\mathscr{O}_U) \xrightarrow{\delta\oplus d} \check{C}^1(\mathscr{O}_X) \oplus \check{C}^0(\Omega_U^1) \xrightarrow{D} \check{C}^1(\Omega_U^1) \oplus \check{C}^0(\Omega_U^2) \xrightarrow{-d\oplus\delta} \check{C}^1(\Omega_U^2)$$

$$D : (\alpha, \beta) \longmapsto (-d\alpha + \delta(\beta), d\beta)$$

the associated total complex. It gives the de Rham cohomology $H^{\bullet}_{\mathrm{dR}}(U)$. Now

$$\begin{aligned}
D : \widetilde{\omega} \longmapsto & \left(\frac{\widetilde{dx}}{y} - \frac{dx}{y}\right) \times \left(d\left(\frac{\widetilde{dx}}{y}\right), \frac{dxdy}{y^2}\right) \\
&= \left(-\frac{(6g_2x^2 - 9g_3x - g_2^2)(g_2'x + g_3')}{y\Delta}dt\right) \times \left(d\left(\frac{\widetilde{dx}}{y}\right), \frac{dxdy}{y^2}\right) \\
&= (Fdt) \times \left(G_1\frac{dtdx}{y}, G_2\frac{dtdx}{y}\right) \\
&\in \check{C}^1(\mathscr{O}_U) \otimes \Omega^1_S \oplus \check{C}^0(\Omega^1_{U/S}) \otimes \Omega^1_S
\end{aligned}$$

where

$$\begin{aligned}
F &:= \frac{(3g_3g_2' - 2g_2g_3')(18x^2 - 3g_2) - \Delta'x - 3g_2g_2'y^2}{6y\Delta} \\
G_1 &:= \frac{-18g_2g_2'x^2 + (9g_3g_2' - 6g_2g_3')x + 2g_2^2g_2' - 9g_3g_3'}{2\Delta} \\
G_2 &:= -\frac{(g_2'x + g_3')}{2y^2}.
\end{aligned}$$

This means

$$\nabla(\omega) = \left((F) \times \left(G_1\frac{dx}{y}, G_2\frac{dx}{y}\right)\right) \otimes dt \in H^1_{\mathrm{dR}}(U/S) \otimes \Omega^1_S.$$

Moreover by using Lemma 6.4 we get

$$\begin{aligned}
\left((F) \times \left(G_1\frac{dx}{y}, G_2\frac{dx}{y}\right)\right) \otimes dt \equiv & \left((F') \times \left(G_1'\frac{dx}{y}, G_2'\frac{dx}{y}\right)\right) \otimes dt \mod \mathrm{Im}\check{C}^0(\mathscr{O}_U) \\
&= \frac{-3(2g_2g_3' - 3g_3g_2')}{\Delta}\omega^* + \frac{\Delta'}{12\Delta} \cdot \omega
\end{aligned}$$

where

$$\begin{aligned}
F' &:= \frac{9g_3g_2' - 6g_2g_3'}{\Delta}\frac{x^2}{y} \\
G_1' &:= \frac{\Delta' + 18(3g_3g_2' - 2g_2g_3')x}{12\Delta} \\
G_2' &:= \frac{y^2\Delta' + 3x(xg_2 + 18g_3)(3g_2'g_3 - 2g_2g_3')}{12y^2\Delta}
\end{aligned}$$

This completes the proof of (6.3).

Next we show (6.4). The proof goes in the same way as above. Let

$$\widetilde{\omega^*} := \left(\frac{x^2}{y}\right) \times \left(\frac{x}{2}\frac{\widetilde{dx}}{y}, \frac{(2g_2x^2 + 3g_3x)dx}{2y^3}\right) \in \check{C}^1(\mathscr{O}_U) \oplus \check{C}^0(\Omega^1_U)$$

be a lifting of ω^*. Then

$$\begin{aligned}
D(\widetilde{\omega^*}) &= \left(-d(\frac{x^2}{y}) + \frac{x}{2}\frac{\widetilde{dx}}{y} - \frac{(2g_2x^2+3g_3x)dx}{2y^3}\right) \times \left(d\left(\frac{x}{2}\frac{\widetilde{dx}}{y}\right), d\left(\frac{(2g_2x^2+3g_3x)dx}{2y^3}\right)\right) \\
&= (F dt) \times (G_1\frac{dx}{y}dt, G_2\frac{dx}{y}dt) \in \check{C}^1(\mathscr{O}_U) \otimes \Omega^1_S \oplus \check{C}^0(\Omega^1_{U/S}) \otimes \Omega^1_S
\end{aligned}$$

where

$$\begin{aligned}
h_1 &= -\frac{2}{3}x^2\Delta' - g_2x(2g_2g_3' - 3g_3g_2') - g_2^3g_2' + 36g_3^2g_2' - 6g_2g_3g_3' \\
h_2 &= -g_2^4g_2'x - 4g_2^3g_3'x^2 - g_2^3g_3g_2' + 27g_2g_3^2g_2'x + 108g_3^2g_3'x^2 + 27g_3^3g_2' \\
F &= \frac{(-6g_2g_2'x + 9g_3g_2' - 6g_2g_3')y^4 + h_1y^2 + h_2}{8y^3\Delta} \\
G_1 &= \frac{-30g_2g_2'x^3 + (4g_2^2g_2' + 9g_3g_3')x + 9x^2(3g_3g_2' - 2g_2g_3') + 2g_2^2g_3'}{4\Delta} \\
G_2 &= -\frac{6g_2g_2'x^3 + 4g_2'x^2y^2 + (9g_3g_2' + 6g_2g_3')x^2 + 6g_3'xy^2 + 9g_3g_3'x}{4y^4}
\end{aligned}$$

By Lemma 6.4 again, we get

$$\begin{aligned}
\left((F) \times (G_1\frac{dx}{y}, G_2\frac{dx}{y})\right) \otimes dt &\equiv \left((F') \times (G_1'\frac{dx}{y}, G_2'\frac{dx}{y})\right) \otimes dt \mod \mathrm{Im}\check{C}^0(\mathscr{O}_U) \\
&= \left(-\frac{\Delta'}{12\Delta}\omega^* + \frac{g_2(2g_2g_3' - 3g_3g_2')}{16\Delta}\omega\right) \otimes dt
\end{aligned}$$

where

$$\begin{aligned}
F' &= -\frac{\Delta'}{12\Delta}\frac{x^2}{y} \\
G_1' &= \frac{-2x\Delta' + 3g_2(2g_2g_3' - 3g_2'g_3)}{48\Delta} \\
G_2' &= \frac{-x(4g_2x + 6g_3)\Delta' + 3(2g_2g_3' - 3g_3g_2')g_2y^2}{48\Delta y^2}
\end{aligned}$$

This completes the proof of (6.4).

6.3 Hodge filtration

Let X and C be smooth projective varieties over R of relative dimension 2 and 1 respectively, and $f : X \to C$ a flat surjective morphism with a section $e : C \to X$ such that the general fiber of f is an elliptic curve. We assume that the following conditions hold.

(Hdg1) There is a closed subscheme $T \subset C$ which is finite etale over R, such that f is smooth over $S := C - T$.

(Hdg2) Put $D := f^{-1}(T)$ and $U := X - D$. Then D_{red} is a relative simple NCD over T.

(Hdg3) The multiplicity of any irreducible component of $D \times_R \overline{x}$ is invertible in R for any geometric points $\overline{x} \to \mathrm{Spec} R$.

$$\begin{tikzcd}
U \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & X \arrow[d, "f"] \arrow[dr, phantom, "\square"] & D \arrow[l] \arrow[d] \\
S \arrow[r] & C & T \arrow[l]
\end{tikzcd}$$

Put

$$H^q_{\mathrm{dR}}((X, D)) := H^q(X, \Omega^\bullet_X(\log D)).$$

Let K is the fraction field of R. Then one has

$$H^q_{\mathrm{dR}}((X, D)) \otimes_R K \xrightarrow{\cong} H^q_{\mathrm{dR}}(U_K/K) := H^q_{\mathrm{zar}}(U_K, \Omega^\bullet_{U_K/K}).$$

Let us denote the Hodge filtration by $F^\bullet H^q_{\mathrm{dR}}((X, D))$:

$$F^p H^q_{\mathrm{dR}}((X, D)) := H^q(X, \Omega^{\bullet \geq p}_X(\log D)) \hookrightarrow H^q(X, \Omega^\bullet_X(\log D)).$$

Define the sheaf $\Omega^1_{X/C}(\log D)$ by the exact sequence

$$0 \longrightarrow f^*\Omega^1_C(\log T) \longrightarrow \Omega^1_X(\log D) \longrightarrow \Omega^1_{X/C}(\log D) \longrightarrow 0.$$

This is a locally free sheaf of rank 1 due to the condition **(Hdg3)**. Put

$$\mathscr{H}_e := R^1 f_* \Omega^\bullet_{X/C}(\log D), \quad \mathscr{H}^{1,0}_e := f_*\Omega^1_{X/C}(\log D), \quad \mathscr{H}^{0,1}_e := R^1 f_* \mathscr{O}_X.$$

As is well-known, the Gauss-Manin connection is extended to $\mathscr{H}_e$:

$$\nabla : \mathscr{H}_e \longrightarrow \mathscr{H}_e \otimes \Omega^1_C(\log T).$$

Theorem 6.6 (cf. [SZ] §5) *There is the natural isomorphism*

$$H^2_{\mathrm{dR}}((X, D)) \cong H^1(C, \mathscr{H}_e \to \mathscr{H}_e \otimes \Omega^1_C(\log D)) \oplus \langle e(C) \rangle_R \tag{6.11}$$

where $\langle e(C) \rangle_R$ denotes the sub R-module generated by the cycle class of $e(C)$. Moreover the Hodge filtration corresponds in the following way.

$$F^1 H^2_{\mathrm{dR}}((X, D)) \cong H^1_{\mathrm{zar}}(C, \mathscr{H}^{1,0}_e \to \mathscr{H}_e \otimes \Omega^1_C(\log T)) \oplus \langle e(C) \rangle_R \tag{6.12}$$

$$F^2 H^2_{\mathrm{dR}}((X, D)) \cong H^0_{\mathrm{zar}}(C, \mathscr{H}^{1,0}_e \otimes \Omega^1_C(\log T)) \tag{6.13}$$

$$\mathrm{Gr}^0_F H^2_{\mathrm{dR}}((X, D)) \cong H^1_{\mathrm{zar}}(C, \mathscr{H}^{0,1}_e) \tag{6.14}$$

Proof. An exact sequence

$$0 \longrightarrow \Omega^{\bullet-1}_{X/C}(\log D) \otimes f^*\Omega^1_C(\log T) \longrightarrow \Omega^\bullet_X(\log D) \longrightarrow \Omega^\bullet_{X/C}(\log D) \longrightarrow 0 \tag{6.15}$$

gives rise to a spectral sequence

$$E_2^{pq} = H^p_{\mathrm{dR}}(C, R^q f_*\Omega^\bullet_{X/C}(\log D)) \Longrightarrow H^{p+q}_{\mathrm{dR}}((X,D)).$$

This yields

$$0 \longrightarrow H^1_{\mathrm{dR}}(C, \mathscr{H}_e) \longrightarrow H^2_{\mathrm{dR}}((X,D)) \longrightarrow H^0_{\mathrm{dR}}(C, R^2 f_*\Omega^\bullet_{X/C}(\log D)) \longrightarrow 0.$$

Since the last term is generated by the cycle class of $e(C)$, we have (6.11). On the other hand (6.15) induces an exact sequence

$$0 \longrightarrow \Omega^{\bullet-1\geq p-1}_{X/C}(\log D) \otimes f^*\Omega^1_C(\log T) \longrightarrow \Omega^{\bullet\geq p}_X(\log D) \longrightarrow \Omega^{\bullet\geq p}_{X/C}(\log D) \longrightarrow 0$$

and this yields

$$\begin{tikzcd}
H^1_{\mathrm{zar}}(C, R^1 f_*\Omega^{\bullet\geq p}_{X/C}(\log D) \to R^1 f_*\Omega^{\bullet-1\geq p-1}_{X/C}(\log D) \otimes \Omega^1_C(\log T)) \arrow[d] & \\
H^2_{\mathrm{zar}}(X, \Omega^{\bullet\geq p}_{X/C}(\log D) \to \Omega^{\bullet-1\geq p-1}_{X/C}(\log D) \otimes f^*\Omega^1_C(\log T)) \arrow[r, "\cong"] & H^2_{\mathrm{zar}}(X, \Omega^{\bullet\geq p}_X(\log D))
\end{tikzcd}$$

Now (6.12), (6.13) and (6.14) easily follow from this. □

Example 6.7 When $C = \mathbb{P}^1_R$, one can compute the Hodge filtration $F^\bullet H^2_{\mathrm{dR}}((X,D))$ in the following way. Let

ϵ_P	0	b	2	3	4	$b+6$	10	9	8
$f^{-1}(P)$	smooth	I_b	II	III	IV	I_b^*	II*	III*	IV*

and put

$$\epsilon := \frac{1}{12}\sum_{P\in C(\overline{K})} \epsilon_P \in \mathbb{Z} \tag{6.16}$$

where K is the fraction field of R and $\overline{K}$ denotes its algebraic closure. The canonical bundle formula says

$$K_X \cong f^*(K_{\mathbb{P}^1} \otimes \mathscr{O}_{\mathbb{P}^1}(\epsilon)) \cong f^*\mathscr{O}_{\mathbb{P}^1}(\epsilon-2), \quad R^1 f_*\mathscr{O}_X \cong \mathscr{O}_{\mathbb{P}^1}(-\epsilon).$$

Let r (resp. s) be the number of additive fibers (resp. multiplicative fibers) in the fibration $f : X_{\overline{K}} \to C_{\overline{K}}$. Then one can show

$$\mathscr{H}_e^{1,0} \cong \mathscr{O}_{\mathbb{P}^1}(\epsilon - r), \quad \mathscr{H}_e^{0,1} \cong \mathscr{O}_{\mathbb{P}^1}(-\epsilon). \tag{6.17}$$

Let us see the image of $F^\bullet H^2_{\mathrm{dR}}((X,D))$ into the in homogeneous de Rham cohomology $H^1_{\mathrm{dR}}(S^o,\mathscr{H})$. By replacing S^o with a smaller open set, we may assume that $U^o := f^{-1}(S^o)$ has a Weierstrass form $y^2 = 4x^3 - g_2x - g_3$. We use the same notations as in the beginning of Appendix. Let t be a coordinate of an affine space $\mathbb{A}^1 \subset \mathbb{P}^1$. Fix $P \in \mathbb{P}^1(R) \setminus S^o(R)$. Put $O_{S^o} = \Gamma(S^o, \mathscr{O}_{S^o})$ and

$$O^m_{S^o} := \mathrm{Im}(\Gamma(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}(mP)) \longrightarrow \Gamma(S^o, \mathscr{O}_{S^o}(mP)) = O_{S^o})$$

where we think of $\mathscr{O}_{\mathbb{P}^1}(mP)$ to be the subsheaf of the constant sheaf of rational functions on $\mathbb{P}^1$. Then there is the natural isomorphism

$$H^1_{\mathrm{dR}}(S^o,\mathscr{H}) \cong \mathrm{Coker}(O_{S^o} \xrightarrow{PF} O_{S^o} \cdot \omega \wedge dt). \tag{6.18}$$

Let $\theta(t)dt$ be a base of the image of the following map

$$R \cong \Gamma(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}((2-r-s)P) \otimes \Omega^1_{\mathbb{P}^1}(\log T)) \longrightarrow \Gamma(S^o, \Omega^1_{S^o}(\log(T \cap S^o))).$$

Moreover let $u(t)\omega$ and $v(t)\omega^*$ be a base of the image of the following maps

$$R \cong \Gamma(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}((r-\epsilon)P) \otimes \mathscr{H}^{1,0}_e) \longrightarrow \Gamma(S^o, \mathscr{H}^{1,0}_e) = O_{S^o}\omega),$$

$$R \cong \Gamma(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}(\epsilon P) \otimes \mathscr{H}^{0,1}_e) \longrightarrow \Gamma(S^o, \mathscr{H}^{0,1}_e) = O_{S^o}\omega^*)$$

respectively. Then, under (6.18), the image of the inclusion $F^2H^2_{\mathrm{dR}}((X,D)) \hookrightarrow H^1_{\mathrm{dR}}(S^o,\mathscr{H})$ is equal to

$$O^{\epsilon+s-2}_{S^o} \cdot u(t)\omega \otimes \theta(t)dt. = O^{\epsilon+s-2}_{S^o} u(t)\theta(t) \cdot \omega \otimes dt.$$

Let us see the image of $F^1H^2_{\mathrm{dR}}((X,D))$. Since $H^1(\mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1}(\epsilon - r)) = 0$, the natural map $\Gamma(\mathbb{P}^1, \mathscr{H}_e \otimes \Omega^1_{\mathbb{P}^1}(\log T)) \to H^1(C, \mathscr{H}^{1,0}_e \to \mathscr{H}_e \otimes \Omega^1_C(\log T))$ is surjective. Let us see the image of

$$\Gamma(\mathbb{P}^1, \mathscr{H}_e \otimes \Omega^1_{\mathbb{P}^1}(\log T)) \longrightarrow H^1_{\mathrm{dR}}(S^o,\mathscr{H}) \cong \mathrm{Coker}(O_{S^o} \xrightarrow{PF} O_{S^o} \cdot \omega \otimes dt). \tag{6.19}$$

To do this, we first note that

$$f(t)v(t)\omega^* \otimes \theta(t)dt - \nabla\left(f(t)v(t)\theta(t)\frac{\Delta}{-3(2g_2g_3' - 3g_3g_2')}\omega\right) \tag{6.20}$$

belongs to $\Gamma(\mathbb{P}^1, \mathscr{H}^{1,0}_e \otimes \Omega^1_{\mathbb{P}^1}(\log T))$ (Prop.6.3). More explicitly

$$\text{(6.20)} = -\left(F(t)\frac{\Delta'}{12\Delta} + F'(t)\right)\omega \otimes dt, \quad F(t) := \frac{f(t)v(t)\theta(t)\Delta}{-3(2g_2g_3' - 3g_3g_2')} \tag{6.21}$$

Therefore the image of (6.19) is generated by (6.21) with $f(t) \in O^{-\epsilon+r+s-2}_{S^o}$ and

$$g(t)u(t)\omega \otimes \theta(t)dt \tag{6.22}$$

with $g(t) \in O^{\epsilon+s-2}_{S^o}$.

References

[AS] Asakura, M. and Sato, K.: *Chern class and Riemann-Roch theorem for cohomology without homotopy invariance.* (preprint).

[CDKL] Chen, X., Doran, C., Kerr, M. and Lewis, J.: *Normal functions, Picard-Fuchs equations and elliptic fibrations on K3 surfaces.* (preprint).

[G] Gillet, H.: *Riemann-Roch theorem for higher Algebraic K-theory.* Adv. Math. 40 (1981), 203–289.

[GL] Gordon, B. and Lewis, J.: *Indecomposable higher Chow cycles on products of elliptic curves.* J. Algebraic Geom. 8 (1999), no. 3, 543–567.

[H] Hartshorne, R.: *On the De Rham cohomology of algebraic varieties.* I.H.E.S. Publ. Math. No. 45 (1975), 5–99.

[R] Ramakrishnan, D.: *Arithmetic of Hilbert-Blumenthal surfaces.* In Number theory (Montreal, Que., 1985), 285–370, CMS Conf. Proc., 7, Amer. Math. Soc., 1987.

[S] P. Schneider: *Introduction to the Beilinson conjectures.* In Beilinson's Conjectures on Special Values of L-Functions (M. Rapoport, N. Schappacher and P. Schneider, ed), Perspectives in Math. Vol.4, 1–35, 1988.

[Sch] Scholl, A.: *Integral elements in K-theory and products of modular curves.* In The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), 467–489, NATO Sci. Ser. C Math. Phys. Sci., 548, (2000).

[Si] Silverman, J.: *Advanced topics in the arithmetic of elliptic curves*. Grad. Texts in Math. 15, New York, Springer 1994.

[AEC] Silverman, J.: *The Arithmetic of elliptic curves*. Grad. Texts in Math. 106, Springer 2009.

[St] Stiller, P.: *The Picard numbers of elliptic surfaces with many symmetries.* Pacific J. Math. **128**, 157–189 (1987).

[SZ] Steenbrink, J. and Zucker, S.: *Variation of mixed Hodge structure. I.* Invent. Math. 80 (1985), no. 3, 489–542.